
Take home exam
Part II

Convex Unconstrained and Constrained Optimization

March 27, 2022

Gloria del Valle Cano
gloria.valle@estudiante.uam.es

Problem 1. (1 point) We have worked out the elementary version of Lagrange multipliers assuming that from $g(x, y) = 0$ we can find a function $y = h(x)$ such that $g(x, h(x)) = 0$. But sometimes what we get is that there is an h such that $g(h(y), y) = 0$. Rewrite the Lagrange multiplier analysis in the lecture slides under this assumption.

For $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$ consider the following minimization problem

$$\min f(x, y) \text{ s.t. } g(x, y) = 0.$$

Assuming the **Implicit Function Theorem** holds, we can find a function $x = h(y)$ s.t. $g(h(y), y) = 0$ and, thus, we can write

$$f(x, y) = f(h(y), y) = \Psi(y).$$

At a minimum y^* with $x^* = h(y^*)$ we thus have

$$0 = \Psi'(y^*) = \frac{\partial g}{\partial x}(x^*, y^*)h'(y^*) + \frac{\partial g}{\partial y}(x^*, y^*). \quad (1)$$

But since $g(h(y), y) = 0$, we also have

$$0 = \frac{\partial g}{\partial x}(x^*, y^*)h'(y^*) + \frac{\partial g}{\partial y}(x^*, y^*) \implies h'(y^*) = -\frac{\frac{\partial g}{\partial y}(x^*, y^*)}{\frac{\partial g}{\partial x}(x^*, y^*)}. \quad (2)$$

Putting together 1 and 2 we arrive at

$$0 = \frac{\partial f}{\partial y}(x^*, y^*)\frac{\partial g}{\partial x}(x^*, y^*) - \frac{\partial f}{\partial x}(x^*, y^*)\frac{\partial g}{\partial y}(x^*, y^*).$$

That is, at (x^*, y^*) , $\nabla f \perp \left(-\frac{\partial g}{\partial y}, \frac{\partial g}{\partial x}\right)$ and, since $\left(-\frac{\partial g}{\partial y}, \frac{\partial g}{\partial x}\right) \perp \nabla g$, we have $\nabla f \parallel \nabla g$ i.e. $\nabla f(x^*, y^*) = -\lambda^* \nabla g(x^*, y^*)$ for some $\lambda^* \neq 0$.

Thus, for the **Lagrangian**

$$\mathcal{L}(x, y, \lambda) = f(x, y) + \lambda g(x, y),$$

we have a minimum (x^*, y^*) there is a $\lambda^* \neq 0$ s.t.

$$\nabla \mathcal{L}(x^*, y^*, \lambda^*) = \nabla f(x^*, y^*) + \lambda^* \nabla g(x^*, y^*) = 0.$$

Problem 2. (3 points) We want to solve the following constrained minimization problem:

$$\begin{aligned} \min \quad & f(x, y) = x^2 + 2xy + 2y^2 - 3x + y \\ \text{s.t.} \quad & x + y = 1, \\ & x \geq 0, y \geq 0. \end{aligned}$$

Argue first that f is convex and then:

- Write its Lagrangian with α, β the multipliers of the inequality constraints.
- Write the KKT conditions.
- Use them to solve the problem. For this consider separately the $(\alpha = \beta = 0)$, $(\alpha > 0, \beta = 0)$, $(\alpha = 0, \beta > 0)$, $(\alpha > 0, \beta > 0)$ cases.

First, we know that f is not convex because it has two convex inequality constraints, and an affine equality one. Also, there are

Problem 3. (1 point) Let $f : S \subset \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex function on the convex set S and we extend it to an $\tilde{f} : \mathbb{R}^d \rightarrow \mathbb{R}$ as:

$$\begin{aligned} \tilde{f}(x) &= f(x) \text{ if } x \in S. \\ &= +\infty \text{ if } x \notin S. \end{aligned}$$

Show that \tilde{f} is a convex function on \mathbb{R}^d . Assume that $a + \infty = \infty$ and that $a \cdot \infty = \infty$ for $a > 0$.

We say that S is a **convex set** if for all $x, x' \in S$ and $\lambda \in [0, 1]$,

$$\lambda x + (1 - \lambda)x' \in S.$$

Let $x, x' \in S$ and $\lambda \in [0, 1]$, so here we cover two cases:

- *First case.* If $x, x' \in S$:

$$\tilde{f}(\lambda x + (1 - \lambda)x') = f(\lambda x + (1 - \lambda)x') \leq \lambda f(x) + (1 - \lambda)f(x') = \lambda \tilde{f}(x) + (1 - \lambda)\tilde{f}(x').$$

Where the first equality holds given that S is convex. The inequality holds because f is convex. And, the last equality raises from the definition of \tilde{f} .

- *Second case.* If $x \notin S$ or $x' \notin S$, we have that

$$\tilde{f}(\lambda x + (1 - \lambda)x') \leq \lambda \tilde{f}(x) + (1 - \lambda)\tilde{f}(x') = +\infty,$$

because $\tilde{f}(y) \leq +\infty, \forall y \in \mathbb{R}^d$.

Since both cases satisfy convexity definition holds, we conclude this function is convex.

Problem 4. (2 points) Prove **Jensen's inequality**: if f is convex on \mathbb{R}^d and $\sum_1^k \lambda_i = 1$, with $0 \leq \lambda_i \leq 1$ we have for any $x_1, \dots, x_k \in \mathbb{R}^d$

$$f\left(\sum_1^k \lambda_i x_i\right) \leq \sum_1^k \lambda_i f(x_i)$$

Hint: just write $\sum_1^k \lambda_i x_i = \lambda_1 x_1 + (1 - \lambda_1) \sum_{i=2}^k \frac{\lambda_i}{1 - \lambda_1} x_i$ for an appropriate v and apply repeatedly the definition of a convex function. Start with $k = 3$ and carry on.

We proceed using an inductive procedure:

- If $k = 1$ then $\lambda = 1$, so we simply have $f(x_1) = f(x_1)$, which is true, and nothing to prove. If $k = 2$ we have the definition of the convexity of f :

$$\lambda_1 + \lambda_2 = 1, \lambda_1, \lambda_2 \geq 0 \implies f(\lambda_1 x_1 + \lambda_2 x_2) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2).$$

- We assume the statement is true for k and consider $k + 1$ points x_1, \dots, x_{k+1} , with coefficients $\lambda_1, \dots, \lambda_{k+1} \geq 0$, $\sum_{i=1}^{k+1} \lambda_i = 1$. The evaluation of the linear combination can be decomposed as

$$f\left(\sum_{i=1}^{k+1} \lambda_i x_i\right) = f\left((1 - \lambda_1) \left(\sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} x_i\right) + \lambda_1 x_1\right).$$

Using this, it is straightforward to use the Jensen's inequality on x_1 and $\sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} x_i$ with coefficients λ_1 and $1 - \lambda_1$ respectively. That is,

$$f\left(\sum_{i=1}^{k+1} \lambda_i x_i\right) \leq (1 - \lambda_1) f\left(\sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} x_i\right) + \lambda_1 f(x_1).$$

We may notice that $\sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} x_i$ verifies the inductive hypothesis, thus,

$$f\left(\sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} x_i\right) \leq \sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} f(x_i).$$

Finally,

$$f\left(\sum_{i=1}^{k+1} \lambda_i x_i\right) \leq (1 - \lambda_1) \sum_{i=2}^{k+1} \frac{\lambda_i}{1 - \lambda_1} f(x_i) + \lambda_1 f(x_1) = \sum_{i=1}^{k+1} \lambda_i f(x_i).$$

Problem 5. (3 points) Prove that the following function is convex

$$\begin{aligned} f(x) &= x^2 - 1, & |x| > 1 \\ &= 0, & |x| \leq 1 \end{aligned}$$

and compute its proximal. Which are the fixed points of this proximal?