

Module 2 Assignment — Computational Lab 2

1. Disease & data choice

Ovarian cancer is a heterogeneous group of malignancies with distinct histologic and molecular subtypes. High-grade serous ovarian cancer (HGSOC) is the dominant subtype and is responsible for a large proportion of ovarian cancer deaths. HGSOC is also characterized by extensive genomic instability and frequent alterations in pathways involved in DNA repair, contributing to its aggressive clinical behavior and making it particularly relevant for multi-omics target discovery (Lisio et al., 2019). These molecular features make ovarian cancer a suitable disease for multi-omics analysis, because integrating transcriptomic, proteomic, and genomic evidence may help identify genes supported across multiple biological layers.

For the transcriptomic layer, I first considered the CPTAC ovarian cancer RNA-seq data from LinkedOmics. However, only tumor RNA data were available, so tumor and normal effect sizes and p-values could not be calculated. So I used Expression Atlas experiment E-GEOD-40595, which provides gene-level log2 fold-change and p-value results for ovarian cancer and normal epithelial cells. The dataset was openly accessible (Papatheodorou et al., 2018).

For the proteomic layer, I used gene-level CPTAC ovarian cancer proteomics data from LinkedOmics. Tumor and normal protein abundance were compared for each gene to obtain a protein effect and p-value. The dataset was openly accessible (Vasaikar et al., 2018).

For the genomic layer, I used gene-level somatic mutation data from the same CPTAC ovarian cancer cohort through LinkedOmics. Mutation frequency was calculated for each gene and used as the genomic association-strength measure. The dataset was also openly accessible (Vasaikar et al., 2018).

The transcriptomic data and the CPTAC proteomic/genomic data came from different cohorts, so they were not matched at the patient level. Therefore, the three omics layers were integrated at the gene level.

2. Top targets

RUNX1T1 was the highest-ranked gene in the multi-omics analysis. Open Targets showed existing associations with ovarian cancer-related disease terms, providing additional support for its biological relevance. This made RUNX1T1 a useful example of a top-ranked gene that is also connected to known cancer biology.(Buniello et al., 2025)

GREB1 also showed ovarian cancer-related evidence in Open Targets, including literature and expression-related evidence. This supports GREB1 as a biologically plausible hit recovered by the multi-omics ranking.(Buniello et al., 2025)

COL3A1 showed some ovarian cancer-related association in Open Targets, but the evidence appeared less established than for the other two genes. For this reason, COL3A1 may be considered a less obvious candidate that could be worth further investigation.(Buniello et al., 2025)

Together, these results suggest that the ranking recovered both biologically supported ovarian cancer-related genes and less obvious candidates that may merit additional study.

3. RNA–Protein Discordance

	gene	rna_log2fc	prot_effect	concordant
0	A2M	-1.6	-1.378499	True
1	A2ML1	0.8	-1.477915	False
2	A4GALT	0.2	-0.409790	False
3	AACS	-0.3	0.120361	False
4	AADAC	-2.9	-0.454076	True

Figure 2. RNA–protein concordance and discordance in ovarian cancer.

A2ML1 showed discordant RNA and protein changes in this analysis, with increased RNA expression but decreased protein abundance.

RNA and protein levels do not always change in the same direction. In ovarian cancer cells, previous proteomic work has shown that protein and mRNA expression changes can be discordant, possibly because of post-transcriptional regulation (Stewart et al., 2006).

Therefore, the opposite RNA and protein changes observed for A2ML1 may reflect regulation after transcription, although the exact mechanism for A2ML1 is not established.

The exact mechanism causing the A2ML1 RNA–protein discordance is not known. In addition, the transcriptomic and proteomic data came from different cohorts, so differences between patient groups may also contribute to the disagreement. Therefore, this result should be interpreted as a gene-level observation rather than a confirmed biological mechanism.

4. Limitations and Interpretation

Since the three datasets were combined by gene rather than by the same patients, this analysis can show which genes have evidence across multiple omics layers. These genes can be considered possible candidates for further study. However, the RNA, protein, and mutation changes may not have happened in the same tumors. Also, a high multi-omics score does not

mean that a gene directly causes ovarian cancer or that it will be a good drug target. More experiments would be needed to confirm these findings.

Reference

- Buniello, A., et al. (2025). Open Targets Platform: Facilitating therapeutic hypotheses building in drug discovery. *Nucleic Acids Research*, 53(D1), D1467–D1475. <https://doi.org/10.1093/nar/gkae1128>
- Lisio, M.-A., Fu, L., Goyeneche, A., Gao, Z.-h., & Telleria, C. (2019). High-Grade Serous Ovarian Cancer: Basic Sciences, Clinical and Therapeutic Standpoints. *International Journal of Molecular Sciences*, 20(4), 952. <https://doi.org/10.3390/ijms20040952>
- Stewart, J. J., White, J. T., Yan, X., Collins, S., Drescher, C. W., Urban, N. D., Hood, L., & Lin, B. (2006). Proteins Associated with Cisplatin Resistance in Ovarian Cancer Cells Identified by Quantitative Proteomic Technology and Integrated with mRNA Expression Levels. *Molecular & Cellular Proteomics*, 5(3), 433–443. <https://doi.org/10.1074/mcp.m500140-mcp200>
- Vasaikar, S. V., Straub, P., Wang, J., & Zhang, B. (2018). LinkedOmics: analyzing multi-omics data within and across 32 cancer types. *Nucleic Acids Research*, 46(D1), D956–D963. <https://doi.org/10.1093/nar/gkx1090>
- Papatheodorou, I., Fonseca, N. A., Keays, M., et al. (2018). Expression Atlas: gene and protein expression across multiple studies and organisms. *Nucleic Acids Research*, 46(D1), D246–D251. <https://doi.org/10.1093/nar/gkx1158>