
Foundations of Data Science

Lecture 7: Model criticism and validation

Prof. Gilles Louppe
g.louppe@uliege.be

LIÈGE université




> *If it (a model) disagrees with experiment, it is wrong. In that simple statement is the key to science.*
>
> — Richard Feynman

Model checking


Newcomb's experiment (1882)

Simon Newcomb measured the speed of light with a rotating mirror. He recorded $N = 66$ measurements of the time light takes to travel a known distance.




Newcomb's 66 measurements, in nanoseconds above 24800 ns.

Example. We model the measurements $\mathbf{x}_{\text{obs}} = (x_1, \ldots, x_N)$ as independent Gaussian draws,



$$\mu \sim \mathcal{U}(-1000, 1000)$$
$$\sigma \sim \mathcal{U}(0.1, 1000)$$
$$x_i \sim \mathcal{N}(\mu, \sigma^2) \quad \text{for } i = 1, \ldots, N.$$

MCMC gives samples $(\mu^{(m)}, \sigma^{(m)})$, $m = 1, \ldots, M$, from the posterior $p(\mu, \sigma \mid \mathbf{x}_{\text{obs}})$.



Posterior samples, with the 2.5%, 50% and 97.5% quantiles.

The posterior mean of μ estimates the passage time, in nanoseconds above 24800 ns,

$$\hat{\mu} = \frac{1}{M} \sum_{m=1}^{M} \mu^{(m)} = 26.199182694915095.$$

Most of these digits are noise: the posterior standard deviation of μ is about 1.4 ns. Report no more digits than the uncertainty supports.

Credible intervals

A credible interval of level $1-\alpha$ is an interval $[a, b]$ such that

$$P(a \leq \mu \leq b \mid \mathbf{x}_{\text{obs}}) = 1 - \alpha.$$

The highest posterior density (HPD) interval is the narrowest one. The equal-tailed interval leaves $\alpha/2$ of the posterior on each side.

From the samples, the 95% equal-tailed interval runs from the 2.5% to the 97.5% quantile of the $\mu^{(m)}$,

$$[\mu_{2.5\%}, \mu_{97.5\%}] = [\mathbf{23.6}, \mathbf{28.9}].$$

The probability is over μ, with the data fixed. A confidence interval instead holds the parameter fixed and describes the variability of the interval over repeated data sets.

Posterior predictive checks


The **posterior predictive distribution** of replicated data $\mathbf{x}^{\text{rep}}$ is

$$p(\mathbf{x}^{\text{rep}} \mid \mathbf{x}_{\text{obs}}) = \int p(\mathbf{x}^{\text{rep}} \mid \theta) p(\theta \mid \mathbf{x}_{\text{obs}}) d\theta.$$

If the model fits, replicated data resemble the observed data. A **posterior predictive check** compares the two.


For Newcomb's data, each replicated data set is drawn in two steps:

- draw $(\mu^{(m)}, \sigma^{(m)}) \sim p(\mu, \sigma \mid \mathbf{x}_{\text{obs}})$;
- draw $x_i^{\text{rep}(m)} \sim \mathcal{N}(\mu^{(m)}, (\sigma^{(m)})^2)$ for $i = 1, \ldots, N$.




The observed data among 19 data sets replicated from the posterior predictive distribution.

To check one aspect of the fit, compare a **test statistic** $T(\mathbf{x}_{\text{obs}})$ with its distribution $p(T(\mathbf{x}^{\text{rep}}) \mid \mathbf{x}_{\text{obs}})$ over replicated data sets.



The mean and the variance of the observed data are typical of the replicated data: the model captures location and spread.



The observed minimum is lower than the minimum of every replicated data set.

The Gaussian model cannot produce the lowest measurements.

The **Bayesian p-value** of a test statistic T is

$$P(T(\mathbf{x}^{\text{rep}}) \geq T(\mathbf{x}_{\text{obs}}) \mid \mathbf{x}_{\text{obs}}) = \int P(T(\mathbf{x}^{\text{rep}}) \geq T(\mathbf{x}_{\text{obs}}) \mid \theta) p(\theta \mid \mathbf{x}_{\text{obs}}) d\theta.$$

A value close to 0 or 1 signals a misfit for T.

A frequentist p-value $P(T(\mathbf{x}^{\text{rep}}) \geq T(\mathbf{x}_{\text{obs}}) \mid \theta)$ conditions on a single value of θ. The Bayesian p-value averages over the posterior.

For Newcomb's data, the Bayesian p-values are

- 0.49 for the mean;
- 0.52 for the variance;
- 1.00 for the minimum.

The model fails on the lower tail of the data.

Residual analysis

Suppose the model is a deterministic function plus additive noise,

$$x_i = f(\theta) + \sigma \epsilon_i,$$

with a scale σ and noise $\epsilon_i \sim p(\epsilon)$. The standardized **residuals** are

$$r_i = \frac{x_i - f(\theta)}{\sigma}, \quad i = 1, \ldots, N,$$

with $\theta \sim p(\theta \mid \mathbf{x}_{\text{obs}})$.


If the model fits, the residuals of the observed data are distributed as the residuals of replicated data. If the posterior is concentrated, both are close to the noise distribution $p(\epsilon)$.

A **quantile-quantile (Q-Q) plot** compares two distributions by drawing the quantiles of one against those of the other. Points on the diagonal mean the two distributions agree.

For the Gaussian model, the standardized residuals are

$$r_i = \frac{x_i - \mu}{\sigma}, \quad i = 1, \ldots, N,$$

with $(\mu, \sigma) \sim p(\mu, \sigma \mid \mathbf{x}_{\text{obs}})$.




Residuals of observed and replicated data (left), and their Q-Q plot (right).
The observed residuals have a heavier lower tail and a narrower core.



Model checking compares the observed data with data simulated from the fitted model, by plots and by test statistics. It reveals **misspecifications**, and points to what the next model must fix.


Model comparison

Example. A Student's t model gives large deviations a larger probability than the Gaussian model:



$$
\begin{aligned}
\mu &\sim \mathcal{U}(-1000, 1000) \\
\sigma &\sim \mathcal{U}(0.1, 1000) \\
\nu &\sim \text{Gamma}(2, 0.1) \\
\epsilon_i &\sim t_\nu \\
x_i &= \mu + \sigma\epsilon_i \quad \text{for } i = 1, \ldots, N.
\end{aligned}
$$

MCMC gives samples from the posterior $p(\mu, \sigma, \nu \mid \mathbf{x}_{\text{obs}})$.



Posterior samples of the t model, with the 2.5%, 50% and 97.5% quantiles.

Bayes factors

Consider two models $\mathcal{M}_1$ and $\mathcal{M}_2$, with prior probabilities $p(\mathcal{M}_1)$ and $p(\mathcal{M}_2)$. Bayes' rule gives the ratio of their posterior probabilities,

$$\frac{p(\mathcal{M}_1 \mid \mathbf{x}_{\text{obs}})}{p(\mathcal{M}_2 \mid \mathbf{x}_{\text{obs}})} = \frac{p(\mathbf{x}_{\text{obs}} \mid \mathcal{M}_1)}{p(\mathbf{x}_{\text{obs}} \mid \mathcal{M}_2)} \frac{p(\mathcal{M}_1)}{p(\mathcal{M}_2)}.$$

The first factor on the right is the **Bayes factor** $\mathrm{BF}_{1,2}$.

The Bayes factor measures how much more probable the data are under one model than under the other. $\mathrm{BF}_{1,2} > 1$ favours $\mathcal{M}_1$.

A common scale reads a Bayes factor

- from 1 to 3 as weak evidence;
- from 3 to 10 as moderate evidence;
- above 10 as strong evidence.

For Newcomb's data, $\log \mathrm{BF}_{t,\mathrm{Gaussian}} \approx 30$: strong evidence for the t model.

The Bayes factor depends on the priors. A prior on μ ten times wider leaves it unchanged, since both models share it. A prior on ν ten times wider, $\mathrm{Gamma}(2, 0.01)$, lowers $\log \mathrm{BF}_{t,\mathrm{Gaussian}}$ to about 25.

A Bayes factor is a ratio of **marginal likelihoods**,

$$p(\mathbf{x}_{\text{obs}} \mid \mathcal{M}) = \int p(\mathbf{x}_{\text{obs}} \mid \theta, \mathcal{M}) p(\theta \mid \mathcal{M}) d\theta,$$

an integral over all the parameters, which rarely has a closed form.

The **Laplace approximation** replaces the posterior by a Gaussian. Let $\hat{\theta} = \arg\max_\theta p(\theta \mid \mathbf{x}_{\text{obs}}, \mathcal{M})$ be the maximum a posteriori (MAP) estimate. Then

$$p(\theta \mid \mathbf{x}_{\text{obs}}, \mathcal{M}) \approx \mathcal{N}(\theta \mid \hat{\theta}, \boldsymbol{\Sigma}),$$

where $\boldsymbol{\Sigma}$ is the inverse Hessian of $-\log p(\theta \mid \mathbf{x}_{\text{obs}}, \mathcal{M})$ at $\hat{\theta}$.

Bayes' rule, evaluated at $\hat{\theta}$, and the approximation give


$$\begin{aligned} p(\mathbf{x}_{\text{obs}} \mid \mathcal{M}) &= \frac{p(\mathbf{x}_{\text{obs}} \mid \hat{\theta}, \mathcal{M}) p(\hat{\theta} \mid \mathcal{M})}{p(\hat{\theta} \mid \mathbf{x}_{\text{obs}}, \mathcal{M})} \\ &\approx p(\mathbf{x}_{\text{obs}} \mid \hat{\theta}, \mathcal{M}) p(\hat{\theta} \mid \mathcal{M}) (2\pi)^{d/2} |\boldsymbol{\Sigma}|^{1/2}, \end{aligned}$$

where d is the number of parameters and $(2\pi)^{-d/2} |\boldsymbol{\Sigma}|^{-1/2}$ is the density of the Gaussian at its mode.



The marginal likelihood averages the likelihood over the prior. A flexible model spreads its prior over many data sets, and gives each a small probability. Comparing marginal likelihoods therefore implements a Bayesian **Occam's razor**: a simpler model is preferred unless the data require a more flexible one.

Example. Polynomial regression of degree 3 and 7, on 20 noisy points.




The marginal likelihood favours degree 3 ($\log p(\mathbf{x}_{\text{obs}} \mid \mathcal{M}) \approx -26$) over degree 7 ($\approx -45$), although degree 7 contains degree 3 as a special case.

Cross-validation

Cross-validation compares models by how well they predict data they were not fitted on.

In k**-fold cross-validation**, the data are split into k folds of N/k observations. Each fold serves once as the validation set, and the other $k-1$ folds as the training set. The scores are averaged over the k splits.



The score of a Bayesian model is the **expected log predictive density** (ELPD),

$$\mathbb{E}_{p_r(\mathbf{x}')}[\log p(\mathbf{x}' \mid \mathbf{x}_{\text{obs}})],$$

where p_r is the data distribution and $p(\mathbf{x}' \mid \mathbf{x}_{\text{obs}})$ is the posterior predictive distribution, fitted on the training set. Cross-validation estimates the expectation with the validation set.

For Newcomb's data, 5-fold cross-validation, summed over the 66 held-out measurements, gives -225 for the t model and -287 for the Gaussian model.

If the posterior is replaced by a point estimate $\hat{\theta}$, the ELPD reduces to the log-likelihood $\log p(\mathbf{x}' \mid \hat{\theta})$ of the held-out data.

The end.