

CS1555 Recitation 11 - Solution

Objective: to practice normalization, canonical forms and finding keys, decomposing relations into BCNF.

Part 1: For each of the following relations R and sets of functional dependencies F, do the following:

- 1) Find the canonical cover (minimal cover) of F.
- 2) Using the canonical cover, find the keys of the R.

1. Consider the following set of functional dependencies F on a relation R (A, B, C, D, E):

$A \rightarrow BC$
 $A \rightarrow D$
 $B \rightarrow C$
 $C \rightarrow D$
 $DE \rightarrow C$
 $BC \rightarrow D$

Finding the canonical form:

- Transform all FDs to canonical form (i.e., one attributes on the right):
 $A \rightarrow B$
 $A \rightarrow C$
 $A \rightarrow D$
 $B \rightarrow C$
 $C \rightarrow D$
 $DE \rightarrow C$
 $BC \rightarrow D$
- Drop extraneous attributes:
B in $BC \rightarrow D$ is extraneous, since we already have $C \rightarrow D$. The set of FDs becomes:
 $A \rightarrow B$
 $A \rightarrow C$
 $A \rightarrow D$
 $B \rightarrow C$
 $C \rightarrow D$
 $DE \rightarrow C$
- Drop redundant FDs:
 $A \rightarrow B$ and $B \rightarrow C$ implies $A \rightarrow C$, so we drop $A \rightarrow C$.
 $A \rightarrow B$, $B \rightarrow C$ and $C \rightarrow D$ implies $A \rightarrow D$, so we drop $A \rightarrow D$.
The set of FDs becomes:

$A \rightarrow B$
 $B \rightarrow C$
 $C \rightarrow D$
 $DE \rightarrow C$

which is the canonical cover of F.

Finding the keys of R:

Observations:

- A and E do not appear in the right hand side of any FDs, so they have to appear in all keys of R.
- $AE^+ : AE \rightarrow AEB$ (since $A \rightarrow B$) $\rightarrow AEBC$ (since $B \rightarrow C$) $\rightarrow AEBCD$ (since $C \rightarrow D$). So AE is a key of R.

In this case, we do not need to consider any other combination, because any other combination containing AE (e.g., AEB) is a super key and not minimal.

2. Consider the following set of functional dependencies F on relation R (A, B, C, D, E, H):

$A \rightarrow C$
 $AC \rightarrow D$
 $E \rightarrow AD$
 $E \rightarrow H$
 $A \rightarrow CD$
 $E \rightarrow AH$

Finding the canonical form:

- Transform all FDs to canonical form (i.e., one attribute on the right):

$A \rightarrow C$
 $AC \rightarrow D$
 $E \rightarrow AD$ becomes $E \rightarrow A$ and $E \rightarrow D$
 $E \rightarrow H$
 $A \rightarrow CD$ becomes $A \rightarrow C$ and $A \rightarrow D$
 $E \rightarrow AH$ becomes $E \rightarrow A$ and $E \rightarrow H$

- Remove redundant dependencies:

$A \rightarrow C$

$AC \rightarrow D$

$E \rightarrow A$

$E \rightarrow D$

$E \rightarrow H$

$A \rightarrow D$

- Drop extraneous attributes:

$AC \rightarrow D$ can be removed because we have $A \rightarrow D$ so C is redundant:

$A \rightarrow C$

$E \rightarrow A$

$E \rightarrow D$

$E \rightarrow H$

$A \rightarrow D$

- Drop redundant FDs: Try removing some dependencies in F and still have a set of dependencies equivalent to F .

$E \rightarrow D$ can be deduced from $E \rightarrow A$ and $A \rightarrow D$ so we can remove $E \rightarrow D$.

The set of FDs becomes:

$A \rightarrow C$

$E \rightarrow A$

$E \rightarrow H$

$A \rightarrow D$

which is the canonical cover of F .

Finding the keys of R :

Observations:

- B and E do not appear in the right hand side of any FDs, so they have to appear in all keys of R .

- BE^+ : $BE \rightarrow AEB$ (because $E \rightarrow A$) $\rightarrow AEBC$ (because $A \rightarrow C$) $\rightarrow AEBCD$ (because $A \rightarrow D$) $\rightarrow AEBCDH$ (because $E \rightarrow H$). So BE is a key of R .

In this case, we do not need to consider any other combination, because any other combination containing BE (e.g., AEB) is a super key and not minimal.

Part 2:

1. Consider the following set of functional dependencies F on relation R (A, B, C, D, E, H):

$A \rightarrow C$

$AC \rightarrow D$

$E \rightarrow AD$

$E \rightarrow H$

$A \rightarrow CD$

$E \rightarrow AH$

The key for R is EB and the following set of functional dependencies constitutes the canonical cover:

$A \rightarrow C, E \rightarrow A, E \rightarrow H, A \rightarrow D$

- a) Using Synthesis Method, construct a set of 3NF relations.

We are starting from a canonical cover. \rightarrow We can skip steps 1, 2 and 3.

Step 4: Group FDs with same determinant:

$A \rightarrow CD$

$E \rightarrow AH$

Step 5: Construct a relation for each group:

$R1(\underline{A}, C, D)$

$R2(\underline{E}, A, H)$

Step 6: If none of the relations contain the key for the original relation, add a relation with the key.

$R3(\underline{E}, B)$

$R1, R2$ and $R3$ are in 3NF and in BCNF.

b) Using Universal Method, decompose R into a set of BCNF relations.

Using $A \rightarrow C$ to decompose R, we get:

$R_1(A, \underline{B}, D, \underline{E}, H)$ in 1NF

$R_2(\underline{A}, C)$ is already in BCNF form

Using $A \rightarrow D$ to decompose R_1 , we get:

$R_{11}(A, \underline{B}, \underline{E}, H)$ in 1NF

$R_{12}(\underline{A}, D)$ is already in BCNF form

Using $E \rightarrow A$ to decompose R_{11} , we get:

$R_{111}(\underline{B}, \underline{E}, H)$ in 1NF

$R_{112}(\underline{E}, A)$ is already in BCNF form

Using $E \rightarrow H$ to decompose R_{111} , we get:

$R_{1111}(\underline{B}, \underline{E})$ in BCNF form

$R_{1112}(\underline{E}, H)$ is already in BCNF form

Group the relations with the same key:

$R_1(\underline{A}, C, D)$

$R_2(\underline{E}, A, H)$

$R_3(\underline{E}, B)$

R_1, R_2 and R_3 are in BCNF form.

2. Consider the following set of functional dependencies F on relation R (A, B, C, D, E):

$A \rightarrow BC$

$A \rightarrow D$

$B \rightarrow C$

$C \rightarrow D$

$DE \rightarrow C$

$BC \rightarrow D$

The key for R is AE and the following set of functional dependencies constitutes the canonical cover:

$A \rightarrow B, B \rightarrow C, C \rightarrow D, DE \rightarrow C$

a) Using Synthesis Method, construct a set of 3NF relations.

We are starting from a canonical cover. \rightarrow We can skip steps 1, 2 and 3.

We can skip Step 4: no FDs with same determinant

Step 5: Construct a relation for each group:

$R1(\underline{A}, B)$

$R2(\underline{B}, C)$

$R3(\underline{C}, D)$

$R4(\underline{D}, \underline{E}, C)$

Step 6: If none of the relations contain the key for the original relation, add a relation with the key.

$R5(\underline{A}, \underline{E})$

$R1, R2, R3, R4,$ and $R5$ are in 3NF and in BCNF.

b) Using Universal Method, decompose R into a set of BCNF relations.

Using $B \rightarrow C$ to decompose R , we get:

$R1(\underline{A}, B, C, D, \underline{E})$ in 1NF

$R2(\underline{B}, C)$ is already in BCNF form

Using $C \rightarrow D$ to decompose $R1$, we get:

$R11(\underline{A}, B, C, D, \underline{E})$ in 1NF

$R12(\underline{C}, D)$ is already in BCNF form

Using $A \rightarrow B$ to decompose $R11$, we get:

$R111(\underline{A}, C, D, \underline{E})$ in 1NF

$R112(\underline{A}, B)$ is already in BCNF form

Using $DE \rightarrow C$ to decompose $R111$, we get:

$R1111(\underline{A}, \underline{E})$ in BCNF form

$R1112(\underline{D}, \underline{E}, C)$ is already in BCNF form

Group the relations with the same key:

$R1(\underline{B}, C)$

$R2(\underline{C}, D)$

$R3(\underline{A}, B)$

$R4(\underline{D}, \underline{E}, C)$

$R5(\underline{A}, \underline{E})$

$R1, R2, R3, R4,$ and $R5$ are in 3NF and in BCNF.