

GABLS1: setup and results

Part II: GABLS1

12/15 completed exports admitted to the scientific comparison. Checked 2026-09-20T01:47:54.123Z.

Grids: 12.5, 6.25, 3.125, 2, and 1 m isotropic; 32^3 , 64^3 , 128^3 , 200^3 , and 400^3 cells, all in the same 400 m cube. Small grids run first.

Each grid has exactly three configurations: WENO9 with no interior closure, WENO5 with no interior closure, and WENO9 with Smagorinsky. Smagorinsky $C_s=0.16$, $Pr=1$; paired seed 123 within each grid. These are controlled Breeze settings, not a reproduction of one named historical LES member.

Forcing: $U_g = 8$ m/s, $V_g = 0$, $f = 1.39e-4/s$. Initial potential temperature 265 K below 100 m and $265+0.01(z-100)$ K above. Initial temperature noise is uniform with 0.1 K peak-to-peak range below 50 m. Surface $\theta_{s,ref}=265-0.25t_{hours}$ K; $\theta_{ref}=263.5$ K is a separate buoyancy reference.

Wall: MOST exchange with $\kappa = 0.4$, $\beta_m = 4.8$, $\beta_h = 7.8$, roughness = 0.1 m for heat and momentum. Impermeable boundaries, free-slip upper lid. Vertical-velocity damping is zero below 300 m and ramps linearly to a 60 s relaxation time at 400 m. MOST uses a minimum wind of 0.01 m/s and caps z/L at 10, with a neutral fallback for nonpositive bulk Richardson number; their activity is diagnosed. Exact implementation, damping and timestep choices are recorded in production provenance.

Diagnostics: 541 instantaneous series records at 60 s intervals; 19 profile times with preceding 30 min means after instantaneous t_0 . Compare 7-8 h and 8-9 h. Final-hour profiles combine bins ending at 30600 and 32400 s; penultimate-hour bins end at 27000 and 28800 s.

Compare mean $u/v/\theta$, jet speed/height/turning, total-stress boundary-layer height, surface heat flux/ u_{star}/L , native-face $w_2/w_3/skewness$, resolved/SGS flux partitions and available TKE-budget terms. Missing SGS energy or budget terms remain unavailable; a residual is not numerical dissipation.

Beare et al. (2006), Boundary-Layer Meteorology 118, 247-272, doi:10.1007/s10546-004-2820-6. Official reduced data archive contains 423 files. Native coordinates and missing values are retained; some NERSC headers give 474-534 min for nominal 8-9 h, so exact window matching is not universal.

GABLS1: setup and results

Cluster agents own implementation, scheduling and audited exports. Desktop monitoring is scheduled every 10 min to collect newly completed cases and rebuild this document. Actual submitted jobs and completed-export evidence are recorded in the status files, separate from the authorized matrix.

Production submission records: 15/15 cases have assigned Slurm jobs. Submitted is distinct from running or scientifically complete. The live queue snapshot below also includes any smoke tests.

Implementation choices: Breeze uses anelastic dynamics and Float32 arithmetic; the reference archive is predominantly Boussinesq. Base pressure is 100000 Pa so the prescribed surface temperature equals surface potential temperature; the corresponding reference density differs from the specification's rounded 1.3223 kg/m³ by about 8 parts per million. Smagorinsky-Lilly uses $C_s=0.16$, $C_b=1$, $Pr=1$.

At $\Delta=12.5$ m, WENO9/Smagorinsky has strongly suppressed resolved turbulence: peak final-hour w_2 is $1.12e-06$ m²/s², $1.21e-05$ times the no-closure value. Maximum magnitudes of the final-hour mean resolved and SGS momentum-flux vectors are $1.34e-07$ and 0.0415 m²/s². Its h is 140.6 m with only 0.242 m standard deviation across final-hour 60 s samples. Similar h or mean wind can therefore coexist with very different resolved turbulence and transport partition; h alone cannot establish fidelity. Skewness in this weak-variance regime must not be interpreted as evidence of vigorous turbulent motions.

At $\Delta=12.5$ m without interior closure, WENO9 versus WENO5 gives final-hour h 195.6 versus 136.1 m and peak resolved w_2 0.0927 versus 0.0574 m²/s². This is a scheme sensitivity at fixed grid and seed, not evidence of resolution convergence.

At $\Delta=6.25$ m without interior closure, WENO9 versus WENO5 gives final-hour h 187.8 versus 150.9 m and peak resolved w_2 0.0976 versus 0.0656 m²/s². This is a scheme sensitivity at fixed grid and seed, not evidence of resolution convergence.

At $\Delta=3.12$ m without interior closure, WENO9 versus WENO5 gives final-hour h 184.4 versus 162.9 m and peak resolved w_2 0.1008 versus 0.0696 m²/s². This is a scheme sensitivity at fixed grid and seed, not evidence of resolution convergence.

At $\Delta=2$ m without interior closure, WENO9 versus WENO5 gives final-hour h 178.5 versus 159.2 m and peak resolved w_2 0.0993 versus 0.0690 m²/s². This is a scheme sensitivity at fixed grid and seed, not evidence of resolution convergence.

GABLS1: setup and results

n032_weno9_none: peak final-hour resolved w_2 ($0.0927 \text{ m}^2/\text{s}^2$) exceeds the largest archived model peak at the same spacing ($0.0602 \text{ m}^2/\text{s}^2$). Model spread is a comparison baseline, not observational truth; stronger resolved turbulence alone does not establish better fidelity.

n064_weno9_none: peak final-hour resolved w_2 ($0.0976 \text{ m}^2/\text{s}^2$) exceeds the largest archived model peak at the same spacing ($0.0779 \text{ m}^2/\text{s}^2$). Model spread is a comparison baseline, not observational truth; stronger resolved turbulence alone does not establish better fidelity.

n128_weno9_none: peak final-hour resolved w_2 ($0.1008 \text{ m}^2/\text{s}^2$) exceeds the largest archived model peak at the same spacing ($0.0904 \text{ m}^2/\text{s}^2$). Model spread is a comparison baseline, not observational truth; stronger resolved turbulence alone does not establish better fidelity.

Resolution comparison for WENO9/none: reducing spacing from 12.5 to 2 m changes final-hour h from 195.6 to 178.5 m and peak resolved w_2 from 0.0927 to $0.0993 \text{ m}^2/\text{s}^2$. Surface heat flux changes from -15.52 to -11.87 W/m^2 . These compare the coarsest and finest completed grids for this configuration; similarity of selected quantities alone is not a convergence or fidelity test.

Resolution comparison for WENO5/none: reducing spacing from 12.5 to 2 m changes final-hour h from 136.1 to 159.2 m and peak resolved w_2 from 0.05744 to $0.06902 \text{ m}^2/\text{s}^2$. Surface heat flux changes from -11.79 to -10.04 W/m^2 . These compare the coarsest and finest completed grids for this configuration; similarity of selected quantities alone is not a convergence or fidelity test.

Resolution comparison for WENO9/smagorinsky: reducing spacing from 12.5 to 2 m changes final-hour h from 140.6 to 188.9 m and peak resolved w_2 from $1.12\text{e-}06$ to $0.09298 \text{ m}^2/\text{s}^2$. Surface heat flux changes from -9.17 to -12.22 W/m^2 . These compare the coarsest and finest completed grids for this configuration; similarity of selected quantities alone is not a convergence or fidelity test.

For WENO9/Smagorinsky, the finest completed grid has $8.3\text{e}+04$ times the coarsest-grid peak final-hour resolved w_2 . At $\Delta=2$ m, maximum magnitudes of final-hour mean resolved and SGS momentum-flux vectors are 0.06499 and $0.04146 \text{ m}^2/\text{s}^2$; these maxima can occur at different heights and should not be treated as fractions of a common total. Final-hour h sample standard deviation is 4.63 m, and mean h changed by 3.31 m between 7–8 and 8–9 h. Read the variance evolution and flux profiles together: this resolution sensitivity does not establish statistical stationarity or identify a unique cause.

GABLS1: setup and results

Final-hour values are descriptive results, not a fidelity ranking. Resolution comparisons must keep the scheme and closure fixed; archive ensemble membership changes with resolution.

The archive reports skewness directly; its averaging convention is not assumed identical to our ratio of time-averaged central moments. No archive third moment is reconstructed from independently averaged variance and skewness. Both Breeze skewness plots mask levels with final-hour variance at or below $1\text{e-}8 \text{ m}^2/\text{s}^2$; the raw diagnostics remain available.

A fixed 1.0 m archive ensemble median (black dotted line) is repeated across resolution panels wherever a matching quantity is available, alongside same-grid archive spread. The 1.0 m archive has two models; pointwise member counts and source files are saved in `reference_data/fixed_1m_medians.json`. Linear totals and resolved TKE are computed per model before taking medians. No matching reference is fabricated for `w3`, time-mean stress magnitude, unmatched budget terms, wall-regime fractions, peak-variance time series or jet time series. The fixed reference aids resolution comparisons but is not an exact solution.

The 1.0 m IMUK archive contains conspicuously large upper-level SGS values, confirmed in the original files: SGS `uw` spans -0.872 to $0.869 \text{ m}^2/\text{s}^2$ and SGS shear production spans -0.283 to $0.793 \text{ m}^2/\text{s}^3$. Tiny values approaching $1\text{e-}100$ abruptly become order-one values above roughly 330 m. This pattern suggests an archive exponent-format problem, but the intended values have not been established. Raw values and resulting median curves are retained, with affected flux/budget panels flagged. Do not interpret those upper-level excursions as a convergence target.

Corrected production (v2) replaces the original campaign after a diagnostic index restriction truncated total-flux/stress profiles and exposed a boundary-layer-height contour-index error. The corrected source preserves the original physics and all three numerical configurations. Expanded CPU checks passed 278 kernel and 426 writer checks; GPU checks passed 268 kernel and 422 writer checks, including every raw profile extent. All 15 cases were resubmitted as arrays 7099-7103, beginning with 32³. Original outputs remain excluded from scientific comparisons and preserved as failure evidence. The collector and single cluster exporter now target v2.

GABLS1: setup and results

The original 400³ WENO9/no-closure and WENO9/Smagorinsky attempts failed during startup and remain excluded. Their fixed initial timestep of 0.5 s implied an advective CFL of 4 for 8 m/s wind and 1 m cells. The separate startup_dt_v1 snapshot changes only initial dt to $\min(0.5, 0.5 \cdot dx/8)$, or 0.0625 s at 1 m. Both variants passed 72 s GPU tests with finite saved bounds: 852/832 iterations and final actual CFL 0.700088/0.700007. The unchanged wizard gave an effective first dt of 0.06875 s. Replacement jobs 7120_1 (WENO9/no closure) and 7120_2 (WENO9/Smagorinsky) were submitted with a one-job array limit; 7120_1 is running and 7120_2 queued at this snapshot. Original WENO5 job 7103_2 continues without duplication. Original failed source/logs/outputs are preserved in attempt history, and replacements require their own job identity, revision and source-hash audits. Passing the short startup tests supports the repair but does not prove full-run stability, uniquely establish the failure cause, or establish that coarser results are independent of startup settings.

n032_weno9_none: final-hour $h = 195.6$ m; $u_{star} = 0.281$ m/s; surface heat flux = -15.52 W/m²; peak mean $w_2 = 0.0927$ m²/s². Between 7–8 and 8–9 h, h changed by -3.7 m and the w_2 profile changed by 0.00438 m²/s² RMS. Simulation wall time 529.7 s (40114 iterations), excluding preceding compilation/startup.

n032_weno5_none: final-hour $h = 136.1$ m; $u_{star} = 0.242$ m/s; surface heat flux = -11.79 W/m²; peak mean $w_2 = 0.05744$ m²/s². Between 7–8 and 8–9 h, h changed by -2.3 m and the w_2 profile changed by 0.00379 m²/s² RMS. Simulation wall time 505.5 s (39025 iterations), excluding preceding compilation/startup.

n032_weno9_smagorinsky: final-hour $h = 140.6$ m; $u_{star} = 0.223$ m/s; surface heat flux = -9.17 W/m²; peak mean $w_2 = 1.12e-06$ m²/s². Between 7–8 and 8–9 h, h changed by 0.1 m and the w_2 profile changed by $4.74e-08$ m²/s² RMS. Simulation wall time 473.4 s (35312 iterations), excluding preceding compilation/startup.

n064_weno9_none: final-hour $h = 187.8$ m; $u_{star} = 0.279$ m/s; surface heat flux = -15.24 W/m²; peak mean $w_2 = 0.09765$ m²/s². Between 7–8 and 8–9 h, h changed by -3.3 m and the w_2 profile changed by 0.00154 m²/s² RMS. Simulation wall time 1050.0 s (80959 iterations), excluding preceding compilation/startup.

n064_weno5_none: final-hour $h = 150.9$ m; $u_{star} = 0.248$ m/s; surface heat flux = -12.01 W/m²; peak mean $w_2 = 0.06558$ m²/s². Between 7–8 and 8–9 h, h changed by -10.9 m and the w_2 profile changed by 0.00248 m²/s² RMS. Simulation wall time 1029.6 s (79613 iterations), excluding preceding compilation/startup.

GABLS1: setup and results

n064_weno9_smagorinsky: final-hour $h = 149.7$ m; $u_{star} = 0.234$ m/s; surface heat flux = -10.16 W/m²; peak mean $w2 = 0.0315$ m²/s². Between 7–8 and 8–9 h, h changed by -21.0 m and the $w2$ profile changed by 0.00986 m²/s² RMS. Simulation wall time 1020.2 s (79350 iterations), excluding preceding compilation/startup.

n128_weno9_none: final-hour $h = 184.4$ m; $u_{star} = 0.270$ m/s; surface heat flux = -13.56 W/m²; peak mean $w2 = 0.1008$ m²/s². Between 7–8 and 8–9 h, h changed by -1.3 m and the $w2$ profile changed by 0.000826 m²/s² RMS. Simulation wall time 4156.8 s (162036 iterations), excluding preceding compilation/startup.

n128_weno5_none: final-hour $h = 162.9$ m; $u_{star} = 0.245$ m/s; surface heat flux = -11.12 W/m²; peak mean $w2 = 0.0696$ m²/s². Between 7–8 and 8–9 h, h changed by 7.9 m and the $w2$ profile changed by 0.00189 m²/s² RMS. Simulation wall time 4058.5 s (160877 iterations), excluding preceding compilation/startup.

n128_weno9_smagorinsky: final-hour $h = 188.1$ m; $u_{star} = 0.277$ m/s; surface heat flux = -13.37 W/m²; peak mean $w2 = 0.08562$ m²/s². Between 7–8 and 8–9 h, h changed by 3.9 m and the $w2$ profile changed by 0.00112 m²/s² RMS. Simulation wall time 4172.6 s (161379 iterations), excluding preceding compilation/startup.

n200_weno9_none: final-hour $h = 178.5$ m; $u_{star} = 0.261$ m/s; surface heat flux = -11.87 W/m²; peak mean $w2 = 0.0993$ m²/s². Between 7–8 and 8–9 h, h changed by 3.3 m and the $w2$ profile changed by 0.000919 m²/s² RMS. Simulation wall time 9396.4 s (253504 iterations), excluding preceding compilation/startup.

n200_weno5_none: final-hour $h = 159.2$ m; $u_{star} = 0.240$ m/s; surface heat flux = -10.04 W/m²; peak mean $w2 = 0.06902$ m²/s². Between 7–8 and 8–9 h, h changed by 1.8 m and the $w2$ profile changed by 0.000677 m²/s² RMS. Simulation wall time 8387.3 s (252093 iterations), excluding preceding compilation/startup.

n200_weno9_smagorinsky: final-hour $h = 188.9$ m; $u_{star} = 0.274$ m/s; surface heat flux = -12.22 W/m²; peak mean $w2 = 0.09298$ m²/s². Between 7–8 and 8–9 h, h changed by 3.3 m and the $w2$ profile changed by 0.00138 m²/s² RMS. Simulation wall time 10660.3 s (252948 iterations), excluding preceding compilation/startup.


Current queue snapshot (job|state|elapsed|start|node or pending reason):

7120_2|PENDING|0:00|N/A|(JobArrayTaskLimit)

7120_1|RUNNING|3:13|2026-09-20T01:44:41|gpu-prod-st-gpu-prod-1


7103_2|RUNNING|2:16:48|2026-09-19T23:31:06|gpu-prod-st-gpu-prod-2

GABLS1 reference archive: final-hour model spread



Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.


GABLS1 reference archive: evolution



Same-grid archive min-max (≥ 2 members)
 1.0 m archive median (fixed reference)
 Same-grid archive median


Archive series interpolated to common times without extrapolation; missing values omitted.
 Membership varies with grid and time. Black dots: the same 1.0 m ensemble median in every row. Heat flux is upward positive.

GABLS1: mean wind and temperature




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.

GABLS1: vertical velocity moments




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.

GABLS1: streamwise momentum transport




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.
 Archive caution: 1.0 m IMUK SGS data contain upper-level anomalies; retained as supplied and flagged in the report.

GABLS1: vertical heat transport




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.
 Archive caution: 1.0 m IMUK SGS data contain upper-level anomalies; retained as supplied and flagged in the report.

GABLS1: turbulence intensity, asymmetry and stress




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.

GABLS1: selected TKE budget terms




Breeze: 8–9 h means; only fully completed and audited exports. Shown budget terms do not form a closed budget. A budget residual must not be interpreted as numerical dissipation; unavailable terms are not replaced by zero. Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled. Archive caution: 1.0 m IMUK SGS data contain upper-level anomalies; retained as supplied and flagged in the report.

GABLS1: crosswind momentum transport




Archive: nominal final-hour profiles, including some NERSC windows of 474–534 min. Breeze curves, when present: 8–9 h means.
 Native heights interpolated without extrapolation; model membership varies with height and grid. The spread is not observational uncertainty.
 Black dots: fixed 1.0 m ensemble median; interpolation precedes the median. Unavailable references are labeled.
 Archive caution: 1.0 m IMUK SGS data contain upper-level anomalies; retained as supplied and flagged in the report.

GABLS1: wall-model regimes and resolved turbulence




Instantaneous 60 s samples. The wall law caps ζ at 10 and uses neutral exchange for nonpositive bulk Richardson number. These fractions identify when wall-model limits participate in the result. No matching archive regime or peak-variance series is available.

GABLS1: resolved vertical-velocity variance




Logarithmic variance axes distinguish weak resolved motions from sustained turbulence. Left: instantaneous profile maxima; right: 8–9 h means. Black dots: fixed 1.0 m median (right only). No archived peak-variance time series. Nonpositive values omitted without a substituted floor.

GABLS1: evolution



Breeze: instantaneous 60 s samples. Archive series interpolated to common times without extrapolation; model membership varies.
 h = height of 5% total surface stress / 0.95; invalid crossings omitted. Black dots: fixed 1.0 m median. Heat flux is upward positive.

GABLS1: low-level jet evolution



Instantaneous 60 s samples of the maximum horizontal-mean wind speed. Direction is atan(v, u) relative to eastward geostrophic wind. Invalid jet diagnostics are omitted; a changing peak height can jump between adjacent grid levels. No archived jet time series is available.