

SurfaceLayerDiffusivity: GABLS1 results

Corrected transport comparison | 21 September 2026 | four matched runs, one seed.

Question: can transport confined to one or two near-wall faces improve coarse LES? Four paired GABLS1 runs use WENO9 at 12.5 m resolution (32^3 cells), a 400 m cube, the same frozen source, and 9 h of integration.

Finding: all three SurfaceLayerDiffusivity variants move the mean u , mean θ and w^2 profiles farther from the fixed 1 m LES median by RMS error over native levels below 200 m. This result applies to this grid and these settings; it is not a general verdict on the closure.

Near-wall response: at 12.5 m, final-hour w^2 falls by 85–93%. Skewness changes from +0.56 in the control to -0.21, -0.25 and -0.42. Domain-integrated resolved TKE falls by 15–20%. The first-face variance is suppressed much more strongly than the integrated energy.

Surface exchange: final-hour u^* rises from 0.279 m/s to 0.292, 0.289 and 0.301 m/s, versus a reference mean of 0.262 m/s. Surface cooling also strengthens, moving farther from the reference. One-face 100 and 300 s responses are similar; two-face support increases local suppression.

Smagorinsky comparison: the earlier 12.5 m WENO9 run ($C_s=0.16$, $Pr=1$) has a final-hour peak w^2 of only $1.12e-6 \text{ m}^2/\text{s}^2$, versus 0.0864 in the control. Its mean- u error is smaller despite almost absent resolved turbulence. Teal curves show this historical context; its source differs from the four matched runs. Log axes reveal the contrast.

Next hypothesis: restrict SLD viscosity to u and v , retaining tracer diffusivity and all other settings. This isolates direct damping of w , but indirect effects through shear and stratification remain. No u/v -only result is included. Single-seed comparisons do not establish uncertainty; the LES median is not observational truth.


Transport split: corrected diagnostics show SGS mixing supplies 86–95% of first-level u -momentum transport in the SLD cases. Total flux magnitude rises by only 6–17%, while its resolved contribution falls sharply. Mean fields and resolved moments exactly reproduce the earlier runs. The new native coefficient–gradient products pass GPU and export consistency checks; this split does not measure WENO numerical mixing. Historical array 7156 fluxes remain excluded; current figures use corrected array 7293.

Reading the figures: native-height profiles are true 8–9 h means; timelines use instantaneous output. Colors, line styles and markers distinguish cases. Exact data, Julia source, reference hashes and diagnostic exclusions accompany this brief; all original DYCOMS/GABLS1 report material is retained.


Surface-layer mixing versus coarse-grid Smagorinsky

GABLS1 · WENO9 · 12.5 m grid (32^3) · final hour, 8–9 h · four matched runs + earlier Smagorinsky


Wind: added near-wall mixing changes the profile





Temperature: smaller changes above the first levels



Vertical variance: strongest suppression near the wall








Third moment: the near-wall sign changes



Teal stars: earlier Smagorinsky source; its w^2 and w^3 nearly coincide with zero on these linear axes.
Black dotted: fixed 1 m LES median. Matched runs use the corrected, GPU-validated flux diagnostics.

Follow the response beyond the mean profile

Surface exchange through 9 h; skewness from the final-hour native-face moments




Wall flux remains usable; resolved TKE excludes SGS energy. Teal: earlier Smagorinsky source.
Skewness omits $w^2 < 10^{-6} \text{ m}^2 \text{s}^{-2}$, masking most Smagorinsky levels; a ratio of tiny moments is not robust.



Smagorinsky nearly eliminates resolved vertical turbulence

12.5 m GABLS1 · WENO9 · log axes expose the difference hidden by linear scales

8–9 h mean variance profile




Instantaneous horizontal variance



Same grid and advection; Smagorinsky is historical context, not a same-revision controlled treatment.
Zeros are omitted on log axes. The left panel averages first; the right panel takes each instantaneous maximum.

SGS transport replaces most of the resolved near-wall flux

GABLS1 · 12.5 m · 8–9 h means · native faces in the lowest 60 m · corrected diagnostics



Colors, line styles and markers identify cases. SGS wall zeros are omitted; total uses the prescribed wall flux. Interior total = resolved + SGS. This split does not quantify WENO numerical mixing. Teal is historical context.

Resolved-flux factor 2: limited response

Matched sensitivity | 21 September 2026 | GABLS1, 12.5 m, WENO9, one interior face, 300 s filter, 9 h, one seed.

Result: crediting twice the resolved transport reduces the added diffusivity, but does not restore resolved turbulence. In the final hour, first-level w^2 rises 7.1%, while peak w^2 falls 6.3% and vertically integrated resolved TKE falls 6.4%. Both factors remain far below the earlier no-closure near-wall variance.

The intervention: factor a multiplies only the signed local filtered resolved flux inside the closure deficit $\max(0, 1 - a F_{\text{resolved}}/F_{\text{target}})$. The same factor is used for momentum and heat. Factor 2 assumes an additional numerical flux equal and aligned with resolved transport. It neither doubles viscosity nor rescales diagnostic fluxes. The experiment does not measure numerical transport or establish that assumption.

Mechanism: final-hour first-face viscosity falls from 1.321 to 1.118 m^2/s (15.4%); heat diffusivity falls from 1.216 to 0.774 m^2/s (36.3%). The first-face SGS share of u-momentum transport remains large: 88.7% versus 86.8%. Reduced closure coefficients therefore produce only a modest change in the resolved/SGS partition.

Moments: at 12.5 m, w^2 changes from 0.008171 to 0.008754 m^2/s^2 , versus 0.06353 without interior closure. The third moment changes from $-1.872\text{e-}4$ to $-1.186\text{e-}4$ m^3/s^3 , and skewness from -0.253 to -0.145. Skewness is the ratio of hourly averaged moments, not an average of instantaneous skewness.

Mean profiles and exchange: u^* falls from 0.2886 to 0.2831 m/s and sensible heat flux changes from -15.54 to -15.38 W/m^2 . Against the fixed 1 m median at native heights $0 < z \leq 200$ m, final-hour RMS errors change by +0.3% for u , -9.2% for θ , and -7.3% for w^2 . Smaller profile errors coexist with lower resolved energy; they do not demonstrate restored turbulence.

Hour sensitivity matters: in 7–8 h, first-level w^2 instead falls 5.1%, integrated TKE falls 10.4%, and the temperature-profile error increases 8.0%. Peak variance and integrated energy decrease in both windows. The near-wall variance increase and temperature improvement in the last hour are not consistent across the two hours. One seed and neighboring hours do not provide an uncertainty estimate.

Comparison integrity: two fresh cases passed strict admission with matching source, grid, seed, initialization and exact 19-profile/541-series schedules. The factor-1 run reproduces every exported profile and time-series value from corrected array 7293's one-face300 case bit for bit. Native physical flux partition, support and coefficients were checked. Fixed 1 m medians are LES references, not observations.


Resolved-flux factor 2: limited response

Execution provenance: both model children completed 32400 s and exited zero. Slurm nevertheless marked the factor-2 batch exit as one. A separate harmless login-shell probe reproduces that pattern through failing `clear_console` in `.bash_logout` under `set-e`; actual batch shell depth was not recorded. The discrepancy and canceled downstream job are preserved. Admission relies on independently complete outputs, durable child exits, matching sources and logs, not a relabeled scheduler success.

Conclusion: this factor-2 test changes coefficients and exchange modestly, but provides no evidence of broad turbulence recovery. It does not identify the numerical flux or establish a calibrated correction factor. All earlier DYCOMS, GABLS1 and SurfaceLayerDiffusivity results remain in the master report. GABLS3's separate humidity-boundary validation is still pending; no new GABLS3 science is included.

Crediting twice the resolved flux: does turbulence recover?


GABLS1 · 12.5 m · WENO9 · one face · 300 s filter · 8–9 h mean



Factor changes the closure deficit for momentum and heat; displayed physical fluxes are unscaled.
Earlier no-closure run is context. This tests an assumed numerical contribution, not a measured numerical flux.

Resolved and SGS transport near the surface

Same final-hour average · native flux faces · physical fluxes remain unscaled



Interior total = resolved + SGS. At the wall, total includes the prescribed flux; artificial SGS wall zeros are omitted.
Each panel includes the fixed 1 m median wherever that reference quantity is available.

Factor 10 mostly switches the closure off

GABLS1 factor-10 sensitivity | 21 September 2026 | 12.5 m, WENO9, one interior face, 300 s filter, 9 h, seed 123. All earlier factor-1/2 results are retained separately.

Result: factor 10 largely removes the added surface-layer mixing after spin-up. Resolved turbulence then resembles the earlier no-closure run. In 8-9 h, first-face w^2 is $0.064671 \text{ m}^2/\text{s}^2$, versus 0.00817056 for factor 1, 0.00875396 for factor 2, and 0.0635343 without closure. Factor 10 is 7.92 times factor 1 and only 1.79% above the no-closure value. This is recovery of resolved turbulence, not proof of accurate total transport.

Mechanism: the multiplier acts only inside the signed local deficit, $\max(0, 1 - a F_{\text{resolved}}/F_{\text{target}})$, for momentum and heat. With $a=10$, aligned resolved transport carrying 10% of the target can satisfy the deficit. Countergradient transport increases it. Crediting ten times the resolved flux assumes a missing contribution nine times as large; the experiment does not measure that contribution. Physical covariance diagnostics are never multiplied.

The saved fractions explain the response. In the final hour, momentum viscosity is exactly zero at 99.9934896% of sampled horizontal point-times; heat diffusivity is zero at 100%. The corresponding raw-zero-deficit and guard-valid-zero-deficit fractions agree. Momentum and heat guards remain valid at 100%, with zero coefficient-cap activity, so this is deficit satisfaction rather than guard failure. Each value averages 60 one-minute samples over all 32×32 horizontal points. It does not imply continuous switch-off between samples: true averaged SGS heat flux is tiny but nonzero, $-1.26136\text{e-}08 \text{ K m/s}$. These six fractions were not saved for factors 1/2 and remain unavailable there.

The closure acts during spin-up. During 0-1 h, mean viscosity is $0.5168 \text{ m}^2/\text{s}$ and heat diffusivity is $0.6870 \text{ m}^2/\text{s}$; zero-coefficient fractions are about 81.54% and 82.98%. By 1-2 h, momentum is zero at 99.9837% of sampled points and heat at 100%. Small intermittent reactivations remain later. In 8-9 h the mean viscosity is $2.72361\text{e-}05 \text{ m}^2/\text{s}$, versus 1.32118 and 1.11797 for factors 1 and 2; sampled mean heat diffusivity is zero.

Moments and energy: factor-10 first-face w^3 is $+0.00977604 \text{ m}^3/\text{s}^3$ and skewness $+0.59443$, compared with negative values for factors 1/2 and $+0.56311$ without closure. Final-hour peak w^2 is 0.0877192 , and integrated resolved TKE is $35.1207 \text{ m}^3/\text{s}^2$, 28.83% above factor 1. The response persists in 7-8 h: first-face w^2 is 0.0668137 , skewness $+0.58543$, and integrated TKE 35.2916 . Neighboring hours and one seed are not an uncertainty estimate.

Factor 10 mostly switches the closure off

Transport and exchange: final-hour SGS carries 0.012852% of first-face u-momentum covariance-plus-SGS transport, versus 88.66% and 86.82% for factors 1/2. Factor-10 u^* is 0.279088 m/s and surface sensible heat flux -15.2328 W/m², close to the no-closure values 0.279164 and -15.2131. The plots' quantity called total is covariance plus actual constitutive SGS. It is NOT the scheme-native WENO advective flux plus SGS: the numerical correction to covariance has not been reconstructed. Agreement of that plotted quantity with a reference cannot validate the missing numerical contribution.


Reference comparison: relative to factor 1, final-hour native-level RMS errors against the fixed 1 m LES median fall 17.33% for u , 9.354% for θ , and 27.43% for w^2 . Nevertheless, factor-10 errors remain larger than the no-closure errors for all three quantities in 8-9 h. The reference is an LES ensemble median, not observations. This strong sensitivity chiefly approaches the no-closure regime; it does not establish factor 10 as a calibrated or faithful correction.

Comparison integrity: factor10 passed separate strict admission (1 admitted, 0 rejected), with 19 native profiles, 541 exact series times, finite values, supported-face SGS and covariance-plus-SGS consistency. Factors 1/2 retain their original admission. Breeze physics commit is identical; factor10's evaluation revision adds read-only diagnostics, with source-specific GPU validation (4,462 checks). Initial native profiles match exactly. Fresh factor1 already reproduces every corrected historical one-face300 profile and series value bit for bit. The earlier no-closure run is labeled as source context. Job7339 used a non-login wrapper and its durable child record verifies exit zero. The preserved older factor2 scheduler discrepancy is documented separately.

Definitions: plotted 8-9 h profiles average the two true half-hour means ending at 30600 and 32400 s; 7-8 h uses 27000 and 28800 s. Series metrics average samples strictly after the hour start through its end. Skewness is the ratio of averaged third moment to averaged variance^(3/2), omitted for $w^2 < 1e-6$. Native model heights are retained; only the fixed 1 m median is interpolated for unweighted RMS errors over $0 < z \leq 200$ m. No WENO flux reconstruction or additional experiment is included.

Factor 10: mean profiles and resolved turbulence


GABLS1 · 12.5 m · WENO9 · one face · 300 s filter · 8–9 h mean



Factor 10 is vermillion, solid, triangles. Reference medians shown where available.
Native heights and physical moments retained; the added numerical-flux credit is an assumption.

Covariance and SGS transport near the surface


Same final-hour average · native flux faces · physical fluxes remain unscaled



Interior total = covariance + actual SGS. This is NOT scheme-native WENO transport: its numerical correction is missing.
Wall total is prescribed; artificial SGS wall zeros omitted. Fixed 1 m medians shown where available.

What the factor changes: coefficients and local switch-off

One supported face at 12.5 m · coefficients from all three factors · switch-off fractions saved only for factor 10



All fraction denominators include every horizontal point. Exact overlap is meaningful; quantitative values are in the report.
Active/guard validity is separate from positive mixing. These six fractions were not saved for factors 1 and 2.