Notes 1

Gabriel Maher

January 29, 2018

 $^{^1\}mathrm{This}$ document contains my notes on miscellaneous topics.

Part I Mathematics

Chapter 1

Linear Algebra

1.1 Matrix Inverse Formulas

This chapter contains formulas for the inverses of various sums of matrices.

1.1.1 Inverse of A + B

Let C + D be the inverse of A + B, then

$$(C+D)(A+B) = I$$

$$\Rightarrow CA + CB + DA + DB = I.$$
(1.1)

Now let $C = A^{-1}$, then

$$A^{-1}A + A^{-1}B + DA + DB = I$$

 $\Rightarrow A^{-1}B + DA + DB = 0.$ (1.2)

Now let $D = A^{-1}MA^{-1}$, then

$$A^{-1}B + A^{-1}M + A^{-1}MA^{-1}B = 0$$

$$\Rightarrow A^{-1} (B + M + MA^{-1}B) = 0$$

$$\Rightarrow B + M + MA^{-1}B = 0.$$
(1.3)

Now let M = BFB, then

$$B + BFB + BFBA^{-1}B = 0$$

$$\Rightarrow B (I + FB + FBA^{-1}B) = 0$$

$$\Rightarrow I + FB + FBA^{-1}B = 0$$

$$\Rightarrow F(B + BA^{-1}B) = -I$$

$$\Rightarrow F = (B + BA^{-1}B)^{-1}.$$
(1.4)

Therefore $D = A^{-1}B(B + BA^{-1}B)^{-1}BA^{-1}$. Substituting for C and D, the inverse of A + B is then

$$(A+B)^{-1} = A^{-1} - A^{-1}B(B+BA^{-1}B)^{-1}BA^{-1}$$
(1.5)

Note that (1.5) only requires A to be invertible. If B is invertible (1.5) can be simplified to

$$(A+B)^{-1} = A^{-1} - A^{-1}B(I+A^{-1}B)^{-1}A^{-1}$$
(1.6)

or

$$(A+B)^{-1} = A^{-1} - A^{-1}(I+BA^{-1})BA^{-1}$$
(1.7)

1.2 Inverse of A + BCD

Using (1.5)

$$(A + BCD)^{-1} = A^{-1} - A^{-1}BCD(BCD + BCDA^{-1}BCD)^{-1}BCDA^{-1}$$

$$= A^{-1} + A^{-1}BCD(BC(D + DA^{-1}BCD))^{-1}BCDA^{-1}$$

$$= A^{-1} + A^{-1}BCD(D + DA^{-1}BCD)DA^{-1}$$

$$= A^{-1} + A^{-1}BCD((D(CD)^{-1} + DA^{-1}B)CD)^{-1}DA^{-1}$$

$$= A^{-1} + A^{-1}B(C^{-1} + DA^{-1}B)^{-1}DA^{-1}$$
(1.8)

1.3 Inverse of AB + CD

Using (1.5)

$$(AB + CD)^{-1} = B^{-1}A^{-1} - (AB)^{-1}CD(CD + CD(AB)^{-1}CD)^{-1}CD(AB)^{-1}$$

$$= B^{-1}A^{-1} - B^{-1}A^{-1}CD(CD(I + (AB)^{-1}CD))^{-1}CDB^{-1}A^{-1}$$

$$= B^{-1}A^{-1} - B^{-1}A^{-1}CD(I + B^{-1}A^{-1}CD)^{-1}B^{-1}A^{-1}$$
(1.9)

Chapter 2

Probability

2.1 Gaussian Distributions

2.1.1 Conditional Multivariate Gaussian Distribution

The PDF of a multivariate Gaussian vector is given by

$$P(\boldsymbol{x}) = \frac{1}{\sqrt{2\pi|\Sigma|}} e^{(\boldsymbol{x}-\boldsymbol{\mu})^T \Sigma^{-1}(\boldsymbol{x}-\boldsymbol{\mu})}.$$
 (2.1)

Let

$$\boldsymbol{x} = \begin{bmatrix} \boldsymbol{x}_1 \\ \boldsymbol{x}_2 \end{bmatrix}, \tag{2.2}$$

and let x_2 be given. We then wish to find $P(x_1|x_2)$, the conditional distribution of x_1 given x_2 . To do this we need to find the block form of Σ^{-1} .

Let

$$\Sigma = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}, \ \Sigma^{-1} = \begin{bmatrix} \tilde{A} & \tilde{B} \\ \tilde{B}^T & \tilde{C} \end{bmatrix}$$
 (2.3)

. Using $\Sigma\Sigma^{-1}=I$ we then end up with the following four equations:

$$A\tilde{A} + B\tilde{B}^T = I, (2.4)$$

$$A\tilde{B} + B\tilde{C} = 0, (2.5)$$

$$B^T \tilde{A} + C \tilde{B}^T = 0, (2.6)$$

$$B^T \tilde{B} + C\tilde{C} = I. (2.7)$$

From (2.5) and (2.6) we obtain

$$\tilde{B} = -A^{-1}B\tilde{C}, \ \tilde{B}^T = -C^{-1}B^T\tilde{A}$$
 (2.8)

Then inserting into (2.4) and (2.7) we get

$$(A - BC^{-1}B^T)\tilde{A} = I \Rightarrow \tilde{A} = (A - BC^{-1}B^T)^{-1}$$
 (2.9)

$$(C - B^{T} A^{-1} B) \tilde{C} = I \Rightarrow \tilde{C} = (C - B^{T} A^{-1} B)^{-1}$$
(2.10)

Using (1.5) we then have

$$\tilde{A} = A^{-1} - A^{-1}B(C + (-B^T)A^{-1}B)^{-1}(-B^T)A^{-1}$$

$$= A^{-1} + A^{-1}B(C - B^TA^{-1}B)^{-1}B^TA^{-1}$$
(2.11)

$$\tilde{C} = C^{-1} - C^{-1}(-B^T)(A + BC^{-1}(-B^T))^{-1}BC^{-1}
= C^{-1} + C^{-1}B^T(A - BC^{-1}B^T)^{-1}BC^{-1}$$
(2.12)

Bibliography