

Representations and the symmetric group - Diaconis data

George McNinch

2024-02-07

character table of S_5

Let's use GAP to find the character table of S_5 .

```
gap> G:=SymmetricGroup(5);
Sym( [ 1 .. 5 ] )
gap> tab:=CharacterTable(G);
CharacterTable( Sym( [ 1 .. 5 ] ) )
gap> Display(tab);
CT1
```

2	3	2	3	1	1	2	.
3	1	1	.	1	1	.	.
5	1	1

	1a	2a	2b	3a	6a	4a	5a
2P	1a	1a	1a	3a	3a	2b	5a
3P	1a	2a	2b	1a	2a	4a	5a
5P	1a	2a	2b	3a	6a	4a	1a

X.1	1	-1	1	1	-1	-1	1
X.2	4	-2	.	1	1	.	-1
X.3	5	-1	1	-1	-1	1	.
X.4	6	.	-2	.	.	.	1
X.5	5	1	1	-1	1	-1	.
X.6	4	2	.	1	-1	.	-1
X.7	1	1	1	1	1	1	1

```
gap>
```

Diaconis example – survey data

This data is taken from the paper (Diaconis 1989)

It describes 5,738 completed ballots rank-ordering 5 candidates.

View a rank-ordered ballot as an element of the symmetric group S_5 ; we want to study the *frequency function* f .

first ranking table

TABLE 2
Percentage of voters ranking candidate i in position j

Candidate	Rank				
	1	2	3	4	5
1	18	26	23	17	15
2	14	19	25	24	18
3	28	17	14	18	23
4	20	17	19	20	23
5	20	21	20	19	20

the regular representation

TABLE 3
Decomposition of the regular representation

M	$=$	V_1	\oplus	V_2	\oplus	V_3	\oplus	V_4	\oplus	V_5	\oplus	V_6	\oplus	V_7
Dim	120	1		16		25		36		25		16		1
SS/120		2286		298		459		78		27		7		0

This diagram shows the decomposition of the regular representation into *isotypic components*.

Be careful: the notation Diaconis is using here does not match that used by GAP above. For example, the representation Diaconis writes as V_3 is the isotypic component determined by the irreducible representation labeled X.5 by GAP.

The second row reflects the decomposition of the frequency function f . Namely, write

$$f = \sum_{i=1}^7 f_i \quad \text{with } f_i \in V_i.$$

The second row entries are the “sums of squares” $\langle f_i, f_i \rangle$.

Remember that we can compute the f_i using the *idempotents* in $\mathbb{C}[G]$.

For example,

$$f_1 = \frac{1}{5!} \sum_{\sigma \in S_5} \sigma \cdot f$$

More generally, if χ_i denotes the character of the irreducible representation L_i with $V_i = \mathbb{C}[G]_{(L_i)}$ then

$$f_i = \frac{1}{5!} \sum_{\sigma \in S_5} \chi_i(\sigma^{-1}) \sigma \cdot f$$

Note that $\langle f_3, f_3 \rangle = 459$ is relatively large (ignoring $\langle f_1, f_1 \rangle$ since f_1 is trivial).

normalizing the first-order data

TABLE 4
First order effects

Candidate	Rank				
	1	2	3	4	5
1	-94	371	165	-145	-296
2	-372	-70	267	268	-92
3	461	-187	-354	-97	178
4	24	-175	-58	16	193
5	-18	62	-19	-41	17

The i, j entry in this table is the number of votes ranking candidate i in the j -th position, minus the sample size over 5. In particular, rows and columns sum to 0.

This normalization can also be achieved as follows:

Let f_2 be the projection on V_2 , and consider the functions

$$\sigma \mapsto \delta_{i, \sigma(j)}.$$

The i, j entry of the preceding table is $\langle f_2, \delta_{i, \sigma(j)} \rangle$

TABLE 5
Second order, unordered effects

Candidate	Rank									
	1,2	1,3	1,4	1,5	2,3	2,4	2,5	3,4	3,5	4,5
1,2	-137	-20	18	140	111	22	4	6	-97	-46
1,3	476	-88	-179	-209	-147	-169	-160	107	128	241
1,4	-189	51	113	24	-9	98	99	-65	23	-146
1,5	-150	57	47	45	43	49	56	-48	-53	-48
2,3	-42	84	19	-61	30	-16	82	-76	-39	72
2,4	157	-20	-43	-25	-93	-76	-56	8	38	112
2,5	22	-44	7	15	-117	69	25	62	99	-138
3,4	-265	-7	72	199	39	140	85	19	-52	-233
3,5	-169	10	88	70	78	44	47	-51	-36	-80
4,5	296	-24	-142	-130	-5	-163	-128	38	-9	267

Interpretation in this last table:

Compute the projection f_3 of f into the component V_3 of $M = \mathbb{C}[S_5]$.

Now, consider the easily understood functions

$$\sigma \mapsto \delta_{\{i, i'\}, \{\sigma(j), \sigma(j')\}}$$

in $\mathbb{C}[S_5]$ for distinct i, i' and distinct j, j' .

The space of these functions is a 100 dimensional subspace of $W \mathbb{C}[G]$.

The entries in the table are the *inner products*

$$\langle f_3, \delta_{\{i, i'\}, \{\sigma(j), \sigma(j')\}} \rangle$$

Summary observations

The data were to elect a president for the American Psychological Association. Candidates 1 and 3 were clinicians while candidates 4 and 5 were academicians, two groups within the association with somewhat divergent perspectives.

In the second-order table, we see a preference for candidates 1 & 3 witnessed by the entry 376 corresponding to the entry for candidates $\{1, 3\}$ and ranks $\{1, 2\}$.

And we see a (slightly smaller) preference for candidates 4 and 5 witnessed by the entry 296 corresponding to the entry for candidates $\{4, 5\}$ and ranks $\{1, 2\}$.

Bibliography

Bibliography

Diaconis, Persi. 1989. "A Generalization of Spectral Analysis with Application to Ranked Data." *The Annals of Statistics* 17 (3): 949–79. <https://www.jstor.org/stable/2241705>.