

# week13-02-curve-fitting

April 7, 2025

## 1 [George McNinch](#) Math 87 - Spring 2025

## 2 Week 13

## 3 Curve-fitting

## 4 Application: Curve-fitting

### 4.1 Linear Regression

Consider a set of data  $y_1, \dots, y_m$  measured at points  $x_1, \dots, x_m$ .

We'd like to find the *best fit* of a straight line through the points  $(x_i, y_i)$  in the plane  $\mathbb{R}^2$ .

Thus, we'd like to find coefficients  $\alpha, \beta$  such that

$$\alpha x_i + \beta \approx y_i \quad (1 \leq i \leq m).$$

We have two unknowns  $a, b$  and  $m$  data points, so if  $m > 2$  this should correspond to an *overdetermined system*.

Application of the method of *Least Squares* discussed above to this problem is known as *linear regression*.

Let's write our system in matrix form, first in the case where  $m = 3$ :

$$\begin{aligned} \alpha x_1 + \beta &= y_1 \\ \alpha x_2 + \beta &= y_2 \\ \alpha x_3 + \beta &= y_3 \end{aligned} \implies \begin{pmatrix} x_1 & 1 \\ x_2 & 1 \\ x_3 & 1 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

Our goal is to find  $a, b$ . Using the method of Least Squares, we consider the normal equations:

$$\text{Write } A = \begin{pmatrix} x_1 & 1 \\ x_2 & 1 \\ x_3 & 1 \end{pmatrix} \text{ and } \mathbf{b} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

So we must solve the matrix equation

$$A^T A \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = A^T \mathbf{b}$$

for  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$

Note that  $A$  has rank 2 provided that  $x_i \neq x_j$  for at least one pair  $i \neq j$ . Thus the matrix  $A^T A$  is invertible, and so there is a unique least-squares solution  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ .

The line  $y = \alpha x + \beta$  is then the *best fit* to the data.

## 4.2 Example:

Let's look at some data of the indicated form.

We first load the arrays  $\mathbf{x}$  and  $\mathbf{y}$ .

```
[1]: import numpy as np
import numpy.linalg as la

x1=np.array( [0.2423001476908293, 0.5131246259108058,
              0.6461774297521172, 1.081718126671276, 1.1066957883683026,
              1.2111192770825507, 1.4107663065070208, 1.4131975649194517,
              1.5419085811341127, 1.54252479268561, 1.6770300002589544,
              1.7551956169308414, 1.7813053516565924, 1.8589163396218567,
              2.1309095689276414, 2.1754282799657187, 2.2037015837896003,
              2.3236943766271545, 2.3350556607482655, 2.4539996228127823,
              2.4576156244809266, 2.923756826621138, 2.9603842447427997,
              2.983549618501935, 3.1807751973214238, 3.2264704664586996,
              3.239574306883238, 3.366259631740789, 3.44332570682743,
              3.499492991470027, 3.5778026905209304, 3.7263844780555875,
              3.748176465962385, 3.7630392531018444, 3.82500728312557,
              3.840703575817898, 3.883434170161646, 3.9204994185997712,
              4.003630198700515, 4.048716471000086, 4.093275391602323,
              4.150511176497722, 4.446436799181704, 4.744462663968328,
              5.006360736646923, 5.061552738339686, 5.186531543235159,
              5.325087321284984, 5.378902191627528, 5.4322553049414335,
              5.579442903857698, 5.657430048039373, 5.676263610166316,
              5.798336865104675, 5.959056467826783, 6.061318417229272,
              6.06964642969886, 6.093223956306707, 6.478158146789935,
              6.536602960843557, 6.659316687948101, 6.687753518577183,
              6.863196862656075, 6.917623848496665, 7.158841237060627,
              7.194629418183499, 7.203946765983015, 7.249367122549896,
              7.261069631339737, 7.287252744675527, 7.346305663767231,
              7.3737179854364925, 7.466367724878489, 7.58659610075152,
              7.715645267370789, 7.782203173483429, 7.785342382089407,
              7.802602022584114, 8.101338980359483, 8.135285482190037,
              8.28839037936026, 8.431493684999166, 8.485586033957622,
              8.85209382371834, 9.047878431454915, 9.05343001103644,
              9.069972112952057, 9.182155314706655, 9.192372350356536,
              9.279577753181126, 9.374682847739509, 9.377466789602028,
```

```

9.399110982574728, 9.480394501454253, 9.566216117340616,
9.575498300284826, 9.586752014027084, 9.672607630688484,
9.682121771410525, 9.966366902650725]])

y1=np.array([10.678513316453799, 9.584529759457341, 8.933389099016741,
8.87862266736914, 9.423027222054186, 7.438013534115137,
6.9311080574968695, 8.290588998022537, 6.9339274272120255,
6.477263126125214, 7.18255726109394, 7.34354559925455,
5.848164724350601, 5.934986418528014, 5.739587800186909,
6.3551105645234145, 4.93116293208016, 5.565267306640516,
6.068819391147042, 4.2422617728095755, 4.339549441040255,
3.936934088821659, 3.036014810980061, 2.899422680159072,
3.288016727330138, 2.483313867446464, 3.620815631303368,
1.7897486715633344, 2.84696838151993, 1.4267338461837338,
2.7080781273543795, 1.1292831820994036, 1.067244502477275,
2.7683112572214843, 2.6058180193619425, 2.648908190074,
1.3990042102813791, 1.1164319243991843, 1.523589585780395,
1.882457911030059, 0.7994184343137196, 0.5503403019975655,
-0.7990859019486943, 0.5249368321797867, -1.1109219031528375,
-2.0949474411711146, -2.638219680107408, -2.5738678800868238,
-2.543579233502919, -2.2101226175665847, -3.586059377975382,
-2.053027871767274, -3.3504551324932588, -3.021599640415171,
-3.3122306023678876, -4.008657108762145, -4.980069709911616,
-3.2827675476924436, -3.895938416357931, -4.874969268341777,
-4.639501130091787, -4.899920594943358, -6.477979961761816,
-7.0497428499160195, -5.571513677706172, -6.787215252732467,
-6.2737864213441, -7.758876101699673, -8.114146089194973,
-6.671329893713388, -6.385436025526582, -6.365364764823218,
-6.303746443776603, -8.130546525025999, -7.723115107002123,
-9.361367822284743, -7.946739978331863, -7.220171496026538,
-9.345989645774747, -8.798860958460038, -9.5706899496564,
-10.362339180159731, -10.595006740315613, -11.585157343328293,
-11.855959857781063, -11.232459127265221, -11.106702212311582,
-11.121677153583464, -11.629010242933319, -12.987414691633676,
-12.918674413364048, -11.073001314425724, -11.797296747359566,
-12.257399767617951, -13.879317776583441, -11.758373608587776,
-13.04069656926525, -13.632789248750406, -12.446490726259036,
-14.854056289273501]])

```

```


[2]: import matplotlib.pyplot as plt

## lists x,y have been populated; lets plot the points
def plot_data(x,y):
    fig, ax = plt.subplots(figsize=(12,6))
    return ax.plot(x,y,"o")

plot_data(x1,y1)

```

[2]: [



Now let's perform "linear regression" by solving the corresponding least squares problem. The main thing to be done is creation of the matrix A.

```
[15]: def lin_fit(x,y):  
    # solve the least squares problem  
    A = np.array([[xx,1] for xx in x])  
    ls=la.lstsq(A,y,rcond=None)  
  
    # get the coefficients and report them  
    alpha,beta = ls[0]  
    print(f"L(x) = {alpha:.03}*x + {beta:.03}")  
  
    # return the linear function determined by these coefficients  
    return lambda x:alpha*x + beta  
  
L = lin_fit(x1,y1)
```

$L(x) = -2.48x + 11.0$

```
[13]: def plot_curve_fit(x0,f,x,y):  
    # graph the line with slope alpha and y-intercept beta, and plot the data  
    # points  
    #  
    fig,ax = plt.subplots(figsize=(12,6))  
    #ax.plot(x,alpha1*x + beta1)  
    ax.plot(x,y,'o')
```

```


ax.plot(x0,f(x0))

return fig,ax

plot_curve_fit(x1,L,x1,y1)

```

[13]: (<Figure size 1200x600 with 1 Axes>, <Axes: >)



### 4.3 Fitting for higher degree polynomials

The preceding example determines a linear function which gives the best fit to the data. But we can also attempt to match other curves.

Let's try the case of a quadratic polynomial. Thus, we are given data  $x_1, x_2, \dots, x_m$  and  $y_1, y_2, \dots, y_m$ , and we wish to find coefficients  $\alpha, \beta, \gamma$  such that

$$\alpha x_i^2 + \beta x_i + \gamma \approx y_i \quad (1 \leq i \leq m)$$

Note that the LHS is the  $i$ -th coefficient in the vector

$$A \cdot \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \quad \text{where} \quad A = \begin{pmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ \vdots & \vdots & \vdots \\ x_m^2 & x_m & 1 \end{pmatrix}.$$

So the coefficients  $\alpha, \beta, \gamma$  of the best-fitting curve

$$Q(x) = \alpha x^2 + \beta x + \gamma$$

are obtained by finding the *least squares* solution for the equation

$$A \cdot \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix}$$

Let's first find the best-fit quadratic polynomial with our previous data set:

```
[5]: #
def quad_fit(x,y):
    # solve the least squares problem
    A = [[xx**2,xx,1] for xx in x]
    res=la.lstsq(A,y,rcond=None)

    # extract & report the coefficients
    alpha,beta,gamma=res[0]
    print(f"Q(x) = {alpha:.03}*x^2 + {beta:.03}*x + {gamma:.03}")


    # return the quadratic function determined by these coefficients
    return lambda x:alpha*x**2 + beta*x + gamma

Q = quad_fit(x1,y1)
```

Q(x) = -0.0103\*x^2 + -2.37\*x + 10.8


```
[6]: plot_curve_fit(x1,Q,x1,y1)
```

[6]: (<Figure size 1200x600 with 1 Axes>, <Axes: >)



```
[7]: plot_curve_fit(np.linspace(-80,60,100),Q,x1,y1)
```

```
[7]: (<Figure size 1200x600 with 1 Axes>, <Axes: >)
```



#### 4.4 New data

We load now some new data x2,y2

```
[8]: x2=np.array([-9.500506098628847, -9.436078956402293,
-9.164729339392766, -8.45233138486573, -8.307739104979152,
-7.828001140674944, -7.705455728811295, -7.571960471086983,
-6.987892876786192, -6.607031656307571, -6.564029048283118,
-6.393814384261753, -5.998800384719204, -5.7183134001218345,
-5.560757660996165, -5.313898241447623, -5.180315921867592,
-4.703049758011512, -4.65291534750819, -4.636925028892335,
-4.473995832831181, -4.349582934094259, -4.276820618215415,
-3.9856919476729056, -3.3024763695520782, -2.930284450264594,
-2.460449933651126, -2.388066289115109, -2.3869772995550846,
-2.382172607813456, -2.282647403270765, -2.2514206929597433,
-2.2151987854220563, -1.8688379031501547, -1.040037196669891,
-0.592982419578636, -0.5285061183473907, -0.07469879485638131,
-0.03672428155128671, 0.08965963140970601, 0.554605101162359,
1.2625261558731822, 1.3951852942052394, 1.422010962689896,
1.5779572321503075, 1.5791882604555276, 1.61640050593763,
1.8955924638126866, 2.4120975394418576, 2.4275213285530626,
2.640546120143096, 3.118535721592007, 3.7246612880274874,
3.778405649802931, 3.818590524157603, 3.8965198726985513,
3.9968635744727568, 4.05853387329822, 4.842259209080241,
```

```

5.2644529452275215, 5.329049050145095, 5.412276483696772,
5.820060572107913, 5.824669115601578, 6.398794721957497,
6.4402471330505975, 6.778030229348008, 6.822865684632326,
7.154884482248647, 7.433033388040794, 7.95313306257955,
8.542338054047299, 8.67214809353717, 8.695899516122513,
8.734604130466039, 8.853944058863625, 8.878828934856866,
9.052318939780356, 9.05661642071038, 9.362282799156798,
9.610149658359706, 10.214633633482066, 10.307112596020925,
10.619306436246774, 10.621964709661611, 10.905709588359596,
11.16622763212822, 11.50521260209901, 11.629057622884542,
11.753805653041166, 11.807760553764172, 12.170454913700738,
12.228946860809256, 12.238348904416945, 12.427719333651723,
13.053306270918945, 14.03578560105586, 14.327026523434942,
14.636621765165128, 14.744885176800103])

```

```

y2=np.array([242.5396610014489, 262.99748345006117,
236.21962830726534, 203.36741916361822, 204.59490171086318,
169.55460040120266, 157.45527545110704, 145.4060333613558,
158.9329865744686, 145.1180030935003, 103.48587998057566,
90.98530811955919, 115.82927769102031, 98.2457769772088,
109.86609808439744, 61.132312437676916, 102.58437854309983,
80.22317148499344, 60.88436606939928, 58.026728933635454,
67.27333189547187, 83.32681082764311, 47.84696391123863,
34.55757018595044, 18.228825041017465, -3.45933210037197,
41.72125747872123, 20.797350275633658, 45.8894100648326,
-1.5174797860946683, 0.636099698366511, 29.059890341434148,
2.311482993391394, -3.4171765808344308, 15.63969965759749,
-12.776680135544654, 23.635189348858646, -18.932585171656093,
31.934447339913138, -7.254615510171753, -7.447589708144465,
21.5170767071797, -10.605455001783469, 37.97690993632591,
-16.28075900816511, 17.514929406199215, -15.41909805330178,
-9.28169529245514, 1.6114757958711792, -2.772531397475941,
31.36303821387554, 48.44461813262329, 21.133120624334147,
55.09416933352425, 46.091773340771034, 32.460057766234485,
51.29572119122864, 84.45347100398384, 100.91738746516664,
118.76834790433631, 89.83233547020102, 103.01856200603174,
127.02275401927923, 126.50985785898953, 106.89953936401625,
138.03228724012408, 154.39572040750687, 167.11136504224805,
166.55137233758427, 185.72301536715477, 184.77395051270676,
252.36187934583052, 226.68621500173575, 207.63736161137572,
251.34665622040202, 229.428513458754, 247.93401134715828,
255.26504804394153, 248.81465414128846, 246.64287358721634,
300.0379742831173, 324.57035644342176, 359.6415758204655,
354.6547519583311, 343.8702990976892, 372.48584664055744,
364.8394942502537, 430.1991994902185, 438.6303667250944,
403.11528216459146, 412.71074061223993, 453.6054297235233,


```



```
478.9207367814926, 481.9332925595611, 450.91801347136175,  
497.7328569627587, 614.3798601360598, 625.8601324671473,  
672.018324952574, 659.0107875019195])
```

```
[9]: plot_data(x2,y2)
```

```
[9]: [<matplotlib.lines.Line2D at 0x7f9c022f3690>]
```




```
[10]: Q2=quad_fit(x2,y2)
```

```
Q(x) = 2.97*x^2 + 1.19*x + 2.21
```

```
[11]: plot_curve_fit(x2,Q2,x2,y2)
```

```
[11]: (<Figure size 1200x600 with 1 Axes>, <Axes: >)
```



```
[12]: L2=lin_fit(x2,y2)
      plot_curve_fit(x2,L2,x2,y2)
```

$$L(x) = 16.4*x + 1.22e+02$$

[12]: (<Figure size 1200x600 with 1 Axes>, <Axes: >)

