


Seabed wreckage PDF under the four 00:19 priority hypotheses (next-run-b, core (b) P(family) mixture)



Grey: seabed wreckage PDF, weighted by settled mass (afloat pieces excluded). Dashed orange: impact PDF, every impact at full weight. Before any seabed-search evidence.
Impacts: end of flight next-run (EoF 3c6319f on core (b) m0011 hand-offs; prior track 289.7°; descent idle floor ON; run by an architecture stand-in), strata free/Davey dynamics+radar/descent-climb/routes mixed at core (b) P(family) 0.6948/0.1527/0.1376/0.0149, held fixed and NOT converged (core split-half 0.71-0.88 < 0.896); seeds equal weight within a stratum. Weights = option_posteriors (EoF 3c6319f).
Seabed field: settling's per-stratum resamples (unchanged recipe), systematically sub-sampled to P(family) proportions, no new draws. Mixture counts free/Davey/descent-climb/routes - none: 200,000/43,955/39,609/4,289; r600_no-offset: 40,000/8,791/7,922/858; both_startup-offset: 40,000/8,791/7,922/858; both_no-offset: 40,000/8,791/7,922/858.
Settling (hypothesis/settling 9823b4e): real ocean, GLORYS12V1 column (TEOS-10) and surface current, ERA5 wind, AusSeabed then GEBCO_2026; breakup table PROVISIONAL; implosion, occupants off.
0.02° grid, Gaussian 0.1° (6 NM), HPD on the smoothed density. Arcs drawn from core reference-289 (core (b)'s own arcs lie ~2 km away). Impacts north of 18° S: not computed, excluded (JSON).
NOT ESTIMABLE: mixed-weight impact ESS below 1,000. Platform: Mac, 2 threads. Stand-in composition script (not module code) around settling's unchanged renderer.
Run labels: core (b) split-half NOT converged; two-tank bookkeeping only; PROVISIONAL-OVERNIGHT; EoF sweep run by a stand-in; settling run by an architecture stand-in, module to review.