


Seabed wreckage PDF under the four 00:19 priority hypotheses (next-run-b, repro-radar stratum)



Grey: seabed wreckage PDF, weighted by settled mass (afloat pieces have no seabed position and are excluded). Dashed orange: impact PDF, every impact at full weight. Before any seabed-search evidence.
Impacts: end of flight next-repro-radar/seed-{1,2,3,4} (next-run-b, repro-radar stratum, prior track 289.7°, source /Users/pete/.claude-science/orgs/9db41e8b-db54-4736-82b9-d77e2a9ad222/workspaces/23b26fb9-3c70-4c8e-909b-ac78b132214a/runs/src-next-repro-radar), seeds pooled with equal weight; weights = option_posteriors (hand-off × 00:19 burst likelihood, × log-on lag density for fuel-exhaustion). Dive class and Boeing-calibrated glide PROVISIONAL-OVERNIGHT. Settling (hypothesis/settling 9823b4e): real ocean, GLORYS12V1 column (TEOS-10 density) and surface current, ERA5 wind, AusSeabed then GEBCO_2026; breakup table PROVISIONAL; implosion and occupants alternatives off. Systematic resampling, one settling draw per resampled impact (200,000 in a; 40,000 in b-d). Impacts north of 18° S lie outside the ocean window: not computed, excluded (counted in the JSON). 0.02° grid, Gaussian 0.1° (6 NM), HPD on the smoothed density.
NOT ESTIMABLE: pooled impact ESS below 1,000; such a panel shows where a few dozen impacts lie, not a posterior. Platform: Mac, 2 threads.
Run labels: core (b) split-half NOT converged; two-tank bookkeeping only; PROVISIONAL-OVERNIGHT; EoF sweep run by a stand-in; settling run by an architecture stand-in, module to review; internal-v1 fuel, one-engine flow 2x source tables (core 06:10).