The Australian sea-floor recordings held cannot exclude any coupling efficiency in the swept range at 95 %





Plain-language note: (a) For impacts in each latitude band of the flight simulation, the chance that all four recorders would have missed the sound, against the share of impact energy that becomes sound. The recorders ran for 5 minutes in every 15 and only two of them cover each arrival, so even a loud impact is missed about 56 % of the time. (b) The upper limit on the coupling efficiency that the absence of a detection allows, with 95 % probability, for each treatment of the 00:19 satellite data; it is almost the same as the limit implied by the assumed range alone. Dive probability 0.5. No detection is claimed.

Technical note: Run J, p = 0.5, alpha = 0.05 (M1b), P_D from block 7 injection-recovery on own backgrounds; P(no detection | eta) = weighted mean over impacts of prod_r (1 - P_D,r) with P_D,r = 0 outside the scored IMOS windows; (b) posterior 95 % quantile under the log-uniform prior on [1e-6, 1e-2] with the non-detection likelihood (filled) and the full likelihood including the amplitude-blind time matches at Portland 3275 (open; gap G11-1); classical 95 % exclusion not reached below 1e-2 in any band. Bands: 1-degree streamed sums over all 313.6 M rows + supplement. PREDICTIVE, NOT CALIBRATED; C1, C2; TL arc grid (IMOS: RAM along the grid, imos_tl_ram_arc.csv, amendment A4). Block 11 prereg 90d7181 + A4 e13a32e; module commits 90d7181, e13a32e; product run-j-gated-v2 (consistent table, built 2026-10-22T05:29:54 local).