
Most probable impact regions with the 00:19 timing value (core run I, not converged)

Dark: 50 % region. Light: 90 % region. Cross: mean position.



(i) The maps show where the impact is most probable. The debris drift and the searches make the region larger, because the searches removed the middle of it. If the Pléiades objects are debris, the region is much smaller and to the north-east. Panel d uses only the free-flight family. The result is not converged: the two halves of the seeds give areas that differ by 13 to 41 %.

(ii) Technical: Core run I: 4 core families x 8 seeds x 100,000 hand-off rows, equal family prior; NOT CONVERGED (split-half waypoints 0.916, floor 0.924; free flight 0.830, descent-climb 0.794, Davey reproduction 0.891). End of flight run I (binary mh370-eof-ri1, 102.4 M impacts): piloted turns and interim pitch-down provisional; no structural limit (G13). 00:19: competing log-on causes q = 1/3, W = 00:10:58-01:15:56, C2 silent, survival factor applied; no treatment uses the R1200 BTO. Ocean: GLORYS12 and GlobCurrent (product-relative windage), equal prior, mixed per row (Bayesian model averaging). Drift h50; unscored rows carried at the mean ratio (<= 0.11 % of weight); Western Australia non-recovery not modelled (G26). Hydroacoustics predictive only, not in these products. Searches: ATSB Phase 2 + Bluefin-21, rho 0.05, point target; Ocean Infinity excluded. Pléiades: rho 0.25, rating 5, six clusters plus four COSMO-SkyMed contacts, conditional on the objects being debris. Strata mixed by core share x 00:19 evidence (ruling C), conditional on equal family prior. Composer pass 2, architecture stand-in. 00:19 R600 BTO Only, competing causes q = 1/3. Regions: 0.1° cells, Gaussian 0.1°, highest density, spherical areas. Panels a-c, e-f: equal-prior core mixture (ruling C); d: free-flight family alone.