


What the chart shows. Each square is a possible impact point; the points are 30 NM apart. The colour shows how well the drift of debris from that point explains the nine identified debris finds. Panels a and b use the same scale. Panel c shows where the two ocean models agree. Panel d shows which model gives the higher value. The GlobCurrent run uses a smaller wind effect, because its surface current already includes some wind drift. The grid now includes added points where the flight model puts impacts. In the far south, few model particles reach the finds, so values there are noisy. The results are interim and are not evidence.

Technical. GLORYS12: production (367 nodes, binary d24060aa8006d3ce) + run A extension (343 nodes, binary e6bb17fe32f22b55; chunks 2-3 run by the architecture stand-in, identical config); GlobCurrent: run A (710 nodes, e6bb17fe32f22b55); Darwin arm64; nodes scored by both models shown (703 of 710); 10^5 particles/node (3 windage classes × 8,334 × 4 ocean-error realisations, seeds 1-4); GLORYS12V1 0.494 m and GlobCurrent my-p1d 0 m, each + ERA5 10 m wind; GSHHG coastline; recovery bandwidth 50 km; node extent reference-289 (prior track 289.7°). Log-likelihood relative to joint maximum (-125.08). c: −|Δ log-likelihood|; outlined where |Δ| ≤ 2σ of combined split-half Monte Carlo noise (σ 1.61 GLORYS12, 1.17 GlobCurrent). Node correlation 0.34. GlobCurrent windage product-relative: c_wind offset drawn per particle U(−0.75%, −0.60%) of U10 (audit F1). 6 GLORYS12 and 1 GlobCurrent nodes unresolved (41.2–39.2°S). 00:19 option not applicable (per-node surface).