





Plain English. No debris was found on Australian coasts. Only coasts where people live can show that debris did not arrive (Pete's ruling, 15 Oct 2026). The drift model gives, for each start point, the share of floating objects that would reach each coast in time. A populated coast is assumed to find and report debris as well as the coasts where debris was found. Two ocean models are used, each with equal weight; the region tracked now reaches 160 E. The populated-coast term is calculated from the measured options with a fitted formula; it is not a separate drift run. Panel a has 0.5 deg steps. The impact PDF is run J (waypoints family), 00:19 data held out. The uncontrolled-descent physics is not calibrated.

Technical. 15-160 E pilot runs on deskstar (17 Oct, drift d51d80a, 178 pilot nodes = every 4th run A node, 1e5 particles per node; GlobCurrent 178/178 scored, left-domain 6.7 %; GLORYS12 177/178, left-domain 12.8 %). Populated = ABS RA 2021 major-city + inner-regional segments (WA Perth-Augusta, SA/Vic/Tas, NSW/Qld). Delta = -k ln(1 + W/(c F)) fitted to measured options B and C per model (GLORYS12 k 5.48, c 0.39; GlobCurrent k 4.60, c 0.37; held-out C residual 0.19 / 0.14 nat, SD 0.62 / 0.49); Delta carried from pilot nodes to run A nodes by inverse-distance-squared (8 nearest). Applied to rebuilt run A surfaces, kernel 50 km; cell 'D0 total q=1/3 (C2 silent)'. Level prior: city/inner ln level 0 relative to the find coasts (option B), post hoc restriction. Gap G26 now modelled as a labelled sensitivity.