
Where the aircraft hit the ocean, for each use of the 00:19 satellite data (before the search evidence)




Note 1: Each panel shows the area that holds 50, 90 and 99 % of the probability of the ocean impact. The panels use the two 00:19 satellite messages in different ways. The heavy black, brown and green lines show sea floor that was searched. The search result is not applied here. The log-on cause mixes fuel exhaustion (weight 3) and other causes (weight 1). The flight model is not yet stable, so the results can change.

Note 2: End of flight on core run J (accepted as a pass by project ruling, 16 Oct 2026): 4 route families with 24/18/16/16 seeds x 100,000 hand-off particles at 00:11; 16 children x 4 descents per particle (waypoints), 8 x 4 (others); family weights 0.9982/0.0013/0.0003/0.0002; binary mh370-eof-rj1 (sha256 19031f41). Round-3 uncontrolled-descent laws; spiral bank limit uniform 52-90 deg; dive to surface p = 0.5. Log-on causes C1 fuel exhaustion (Erlang lag) + C2 other (1/W, W = 00:10:58-01:15:56 UTC, silent), q = 1/3; survival factor (no in-flight breakup before 00:19:37) applied; transmission at 00:19 imposed. No treatment uses the 00:19:37 range. Treatments: D0' held out, D1 R600 BTO, D2 R600 BTO+BFO, D3 both BFOs raw, D4 both BFOs + shared start-up offset (Holland), D5 R600 BFO raw + R1200 BFO with offset U[17,130] Hz. Families: A1 = code 1, A2 = code 2, B = codes 3+6 (B uncontrolled about 0.001). PROVISIONAL: piloted-turn model, INTERIM pitch-down. Panel d = equal-prior average of D3/D4/D5 by ln Z (posterior weights 0.381/0.408/0.212). Density 0.02 deg grid, Gaussian sigma 0.08 deg, HPD 50/90/99 %. Searched ground: coverage >= 0.5 outlines of the seabed-search rasters. ESS = Kish effective impacts pooled over seeds. Masses reproduce table-0019-mixture.json (ln Z and ESS identical). Map 86.5-104 E, 40.2-25 S; probability outside it is printed per panel (north of 30 S: 1.6 % / 1.5 % / 1.4 % / 1.3 %).