

Simple words: This chart uses the satellite data to 00:11 and our model of the last part of the flight. It adds a second set of flight paths that goes more to the north. We add the two sets by a fixed rule. The rule does not change the answer; it only makes the north less noisy. The physics of the last part of the flight is not yet calibrated.

Technical: source posterior = core run J, waypoints stratum (24 seeds) + run-J northern supplement (16 seeds, every 2nd row, multiple importance sampling by the balance heuristic with r(00:11 latitude) from r_hat.csv); prior track = run J waypoint route sampler; 00:19 data option as labelled, log-on cause q = 1/3 with 'other cause' silent, survival factor applied; provisional physics: round-3 descent laws, bank limit U[52, 90] deg, interim pitch-down, uncontrolled arm uncalibrated (Boeing gate failed); no search, ocean-drift or other evidence applied (impact level); waypoints stratum only (core prior share 0.998). ESS = Kish effective sample size over distinct 00:11 states (identical resampled copies merged). Grey bars: core's predicted ranges posted before the run (P5); a bar without a top is a lower bound only; they apply to the 64-descent set. Extra descents: 256 more per 00:11 path north of 35.5 S, pooled with the first 64 per path by count, for the both-burst options only. The last option is scored with a 4.3177 Hz frequency error, as run J.