

Plain: The debris model gives a number for how well each impact place explains the debris that was found (flaperon and eight other items, with the dates they were found) and the absence of finds on Australian coasts where people live. We compare the impact places that agree with the sound from H01W (bearing 306.18 deg; arrival 00:52:03 or 00:54:30) with all impacts from the flight-path calculation. A positive number means the debris favours the place near the bearing. The two ocean models have equal weight. The selected impacts are few, so the effective sample size is small.

Technical: drift surfaces run A (rebuilt, h50 kernel, GLORYS12V1 + ERA5 and GlobCurrent + ERA5, equal-weight likelihood mixture), Australian populated-coast term reconstructed with the saturating law from the 15-160 E pilot (k,c fitted per model), release 00:19:37 UTC in these surfaces (dedicated L-point run at the branch impact times queued). Prior: run J four families (waypoints 0.9982, free, repro-radar, descent-climb), cell 'D{d} total q=0.333 (C2 silent)', 74 seeds. Acoustic weight: pre-registered Q2 likelihood, c = 1.48 km/s, s_b = 0.5 deg (core). Bars: 2-sigma = transport noise (100 mean-preserving node perturbations per 1-deg band split-half SD) and seed bootstrap (100) combined. Uncontrolled-descent physics not calibrated. No search-area term (separate module); joined planned descents J0011/J2241 not included. Lines in (a): run J median impact longitude per 0.5 deg latitude (arc-latitude first look). Rows outside drift support (2-5 % of weight) carried at the mean.