

(i) Plain: The data is from the run J flight model (simulated flights after 00:11, end-of-flight descents). We use only flights that agree with the satellite data. We give each flight a weight for how well its impact time and place agree with the sound that Kadri (2024) found at the H01W hydrophone on bearing 306.18 deg. Shaded areas contain 50 %, 90 % and 99 % of the weighted impacts. The sample near the bearing is small, thus the areas are not stable. Haixun 01 is a secondary check only. Searches, drift and other evidence are not used.

(ii) Technical: source posterior end-of-flight run J (strata waypoints/free/repro-radar/descent-climb, P(family) 0.9982/0.0013/0.0003/0.0002, 74 seeds), cells "D{d} total q=0.333 (C2 silent)" (log-on cause mixture p(C1) = 0.75, survival factor applied); physics PROVISIONAL (interim pitch-down, piloted turns, residual roll P(left) 0.8, round-3 free-flight laws); acoustic likelihood per pre-registration Q2 (c = 1.48 km/s, s_t^2 = (d 0.015/1.48^2)^2 + 25 s^2, s_b = 0.5 deg); grid 0.05 deg; HPD 50/90/99 %; search case: none; ocean model: none; exclusions: none; Kish ESS < 1,000 in every panel, so every panel is NOT ESTIMABLE (end-of-flight map threshold).