
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 R600 BTO + Raw BFO drift-error correlation 0.25, one ocean model (GLORYS12) — core run C NOT CONVERGED



	ρ = 0.25 GLORYS12 (drawn)	ρ = 0 GLORYS12	ρ = 0.5 GLORYS12	ρ = 0.25 two ocean models*
50 % region, km²	5,817	5,470	6,138	7,582
90 % region, km²	30,998 σ not run	28,478 ± 214	33,493 σ not run	41,597 σ not run
mean position	36.01 S 92.38 E	35.96 S 92.38 E	36.05 S 92.37 E	35.95 S 92.36 E
outside past searches	74.3 %	73.0 %	75.0 %	69.9 %
share left by searches	0.412	0.395	0.421	0.366
tension ln S (p)	+0.13 (0.43)	-0.01 (0.38)	+0.24 (0.47)	-0.32 (0.30)
mean shift from PDF without H	36 NM	38 NM	34 NM	38 NM
PDF without H inside 90 % region	15 %	14 %	15 %	16 %

- probability density if the objects are debris (relative)
- 50 % region (drawn setting)
- 90 % region (drawn setting)
- 90 % region, drift-error correlation 0, GLORYS12
- 90 % region, drift-error correlation 0.5, GLORYS12
- 90 % region, drift-error correlation 0.25, two ocean models*
- impact PDF without H (sampled points)
- ATSB Phase 2 2014-17 (official)
- Ocean Infinity 2018 (grade C)
- OI 2025-26 south-east band (grade C)
- OI north-west band (not searched; grade C)
- 7th arc, FL400
- Pléiades rating-5 objects
- COSMO-SkyMed radar contacts (all four)

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses one ocean model (GLORYS12) and a correlation of 0.25 between the drift errors of the two satellite data sets. The thin lines show the 90 % region for other settings. *The two-model result counts the wind drift of the GlobCurrent model two times, thus it is shown only for comparison. The flight calculation (core run C) is not converged. Use these results with caution.

Technical: core run C (4 strata x 8 seeds) split-half NOT CONVERGED (floor 0.924: free 0.918, Davey dynamics + radar 0.910, descent-climb 0.892, routes 0.850); end-of-flight physics PROVISIONAL (descents > 6,500 ft/min and unloading not reachable; no structural limit, G13). 00:19 R600 BTO + Raw BFO, existence constraint 'unpowered'; strata re-weighted by the 00:19 evidence (family-evidence-run-c.json, key '00:19 R600 BTO + Raw BFO +unpowered': next-c-free 0.606, next-c-repro-radar 0.218, next-c-descent-climb 0.154, next-c-routes 0.022). After Phase 2 + Bluefin-21 (official) + OI 2018 + OI 2025-26 SE band (grade C; NW band drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors per component (prepare/rho_surface.py), ρ = 0.25 headline (ocean transport measured 0.07-0.41, GLORYS12), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 only (architecture ruling 2, 11 Oct 02:15 -0600); *both models equal weight, GlobCurrent windage double-counted (drift audit F1). ± = split-half σ (rms over three 4 + 4 seed partitions), from the module's splithalf.csv; measured for ρ = 0 GLORYS12 only. The σ for the other three settings was not run (the heavy-compute lock was held by other work). Conditional on H; no Bayes factor or P(H | data). Holland H1/H2 not estimable. Background: 6,000 points from the impact PDF without H after the same searches. Stand-in redraw (module scripts, inputs changed only); module review pending.