
If the imaged objects are from 9M-MRO: where the debris entered the sea — 00:19 data held out, transmission imposed (D0′) core run J, drift-error correlation 0.25, two ocean models (headline)



	ρ = 0.25 two models (drawn)	ρ = 0.25 GLORYS12 only	ρ = 0 two models	ρ = 0.5 two models
50 % region, km²	11,870	11,760	10,079	13,554
90 % region, km²	63,129	56,633	50,932	72,534
mean position	35.44 S 92.20 E	35.44 S 92.25 E	35.39 S 92.25 E	35.49 S 92.15 E
outside official search outlines	37.0 %	37.9 %	36.4 %	36.9 %
tension ln S (p)	-1.04 (0.22)	-1.02 (0.23)	-1.02 (0.23)	-1.03 (0.22)
mean shift from PDF without H	109 NM	110 NM	112 NM	105 NM
PDF without H inside 90 % region	33 %	15 %	22 %	40 %

- probability density if the objects are debris (relative)
- 50 % region (drawn setting)
- 90 % region (drawn setting)
- 90 % region, drift-error correlation 0.25, GLORYS12 only
- 90 % region, drift-error correlation 0, two ocean models
- 90 % region, drift-error correlation 0.5, two ocean models
- impact PDF without H (sampled points)
- ATSB Phase 2 2014-17 (official)
- Ocean Infinity 2018 (grade C) — drawn only, not applied
- OI 2025-26 south-east band (grade C) — drawn only, not applied
- OI north-west band (not searched; grade C)
- 7th arc, FL400
- Pléiades rating-5 objects
- COSMO-SkyMed radar contacts (all four)

STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The blue colour shows the probability with this assumption. The grey points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets; the thin lines show the 90 % region for other settings. No seabed search is applied. The flight calculation is core run J; its sampling is accepted as converged by project ruling. The result is conditional on this assumption only.

Technical: core run J (waypoints 24, free 18, descent-climb 16, repro-radar 16 seeds; strata by core P x own evidence; D3 and D4 waypoints from the both-burst supplementary set, 8 seeds); split-half accepted as a pass by project ruling, 16 Oct 2026. End-of-flight physics PROVISIONAL (piloted-turn model, INTERIM pitch-down, push-over and Boeing sequence; end of flight's run J labels). 00:19 data held out, transmission imposed (D0′); log-on cause: competing-causes total, q = 1/3, C2 silent (primary); survival factor applied; end of flight's own row weights (option_table_0019.seed_terms). Search: none applied (outlines drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors, ρ = 0.25 (measured 0.07-0.41), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight, GlobCurrent windage product-relative (drift audit F1). Headline (brief v2); GLORYS12 only beside. Maps from per-impact columns on the wide 0.05 deg grid (prepare/runj_closeups.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data). Coverage: effective impacts 141,104 under H (1,946,710 without H); A1 under H 56,354. Gate (1,000) met. Background: 6,000 points from the impact PDF without H, before any seabed search.