
If the imaged objects are from 9M-MRO: where the debris entered the sea, on GEBCO 2026 relief — 00:19 R600 BTO + R600 BFO — core run J



STE100: This map shows where the aircraft possibly went into the sea, if the objects in the Pléiades satellite images are from 9M-MRO. The red bands show the 50, 90 and 99 % regions with this assumption. The points show the probability without this assumption. The drawn result uses two ocean models with equal weight and a correlation of 0.25 between the drift errors of the two satellite data sets. No seabed search is applied. The flight calculation is core run J; its sampling is accepted as converged by project ruling. The result is conditional on this assumption only.

Technical: core run J (waypoints 24, free 18, descent-climb 16, repro-radar 16 seeds; strata by core P x own evidence; D3 and D4 waypoints from the both-burst supplementary set, 8 seeds); split-half accepted as a pass by project ruling, 16 Oct 2026. End-of-flight physics PROVISIONAL (piloted-turn model, INTERIM pitch-down, push-over and Boeing sequence; end of flight's run J labels). 00:19 R600 BTO + R600 BFO; log-on cause: competing-causes total, q = 1/3, C2 silent (primary); survival factor applied; end of flight's own row weights (option_table_0019.seed_terms). Search: none applied (outlines drawn only). Under H: 12 rating-5 Pléiades objects (6 clusters, 3 km, equal weight) x all four COSMO-SkyMed contacts, one debris field; bivariate-normal transport errors, ρ = 0.25 (measured 0.07-0.41), ρ = 0 / 0.5 beside; GLORYS12 + ERA5 and GlobCurrent daily + ERA5, equal weight, GlobCurrent windage product-relative (drift audit F1). Headline (brief v2); GLORYS12 only beside. Maps from per-impact columns on the wide 0.05 deg grid (prepare/runj_closeups.py). Conditional on H; no identity likelihood, no Bayes factor or P(H | data).

Coverage: effective impacts 9,953 under H (92,216 without H); A1 under H 3,311. Gate (1,000) met.

Result, before any seabed search: 90 % region 39,188 km², mean 35.52 S 92.34 E; tension ln S -1.23 (p 0.20), mean shift 116 NM; PDF without H inside the 90 % region 34 %. Beside it, GLORYS12 only at ρ = 0.25: 35,978 km², 35.49 S 92.40 E, ln S -1.22.

Relief: GEBCO 2026 15-arcsecond grid (doi:10.5285/4f68d5c7-45eb-f999-e063-7086abc036fa), shown at 0.025 deg; contours every 1 km depth. Right: probability per degree of latitude over the whole grid, before any seabed search.

Background: the impact PDF without H (flight posterior alone, before any seabed search), as 6,000 points drawn in proportion to its probability from 0.05 deg grid cells and jittered within each cell. 89 % of it lies inside this frame.