Published-route result for 18:22 to 19:41: audit of the turn-south gain (UNCONVERGED)



STE100: (a) The bars show how much better a published route (west to ANOKO, then south to ISBIX) predicts each measurement than the model with random turns. Grey bars are radar times. Blue bars are satellite times. (b) We moved the turn point along the westbound track and did the calculation again. The best turn point is 10 NM east of ANOKO. There, the aircraft flies on the published leg. The free-turn model also has these paths, but with less prior probability. The results are not converged. Use them only to compare.

Technical: core early-flight evidence (no end-of-flight or impact modules), 18:01:49 radar prior, satellite BTO/BFO 18:25 to 00:19 (00:19a/b as in runs C/D), radar 18:04 to 18:22 in the likelihood; run C fuel stack; Davey (2016) dynamics, 15 deg bank; mode prior 1/5. (a) per-epoch increments of the 5-mode mixture log-evidence, mean of seeds: pilot route 11 with turn_backs (200k, 2 seeds, ad50) minus run D free (7M, 4 seeds). (b) audit smoke, 50k, seeds 1-2, pseudo-fix on the NILAM-ANOKO geodesic then ISBIX, turn_backs, binary 861e4408-dirty; run D free -135.90; routes family (48 routes, log-mean-exp) -131.59. Split-half not converged; numbers provisional.