

Simplified English: A1 is a cruise to fuel exhaustion, then no control. A2 is a cruise to fuel exhaustion, then a controlled descent. B is a planned descent with a ditching approach that starts before fuel exhaustion. The main run J samples had too few A1 flights when both 00:19 frequency values are used. An extra flight-end run sampled only the most important A1 case. We used its results in place of that part of the main samples, with a known scale factor. Now A1 has enough samples for every use of the 00:19 data (panel a). The search result for A1 and for all types together does not change by more than the sampling noise (b, c). The level of detection in each searched area changes the result much more than the samples do (d). The Ocean Infinity search areas come from community tracings and are a best estimate. Some parts of the flight-end physics are provisional.

Technical: Z = E[P(no find)] under the end-of-flight posterior; P = rho + (1 - rho) prod_k (1 - q_k f_k c_k), base = ATSB Phase 2 (q 0.945) + Bluefin-21 (0.9) + Ocean Infinity 2018 (0.9, f 0.889) + Ocean Infinity 2025-26 outboard band (0.9, f 0.7808), rho 0.05: Ocean Infinity coverage inferred (grade C), included as best estimate; planned next Ocean Infinity area not evidence. Source posterior: core run J (split-half accepted as a pass by project ruling, 16 Oct 2026), prior track = waypoint routes stratum only (core share 0.9982); end of flight run J production (24 seeds, 153.6 M rows) + A1-branch conditional pass (onset flame-out-associated, no intervention; 8 seeds, 13.4 M rows, seeds 2-8 Horvitz-Thompson thinned), stratified estimator, settling's single measure: pass mass x 0.14108, production branch rows 0, P_D0'(branch) = 0.0935. Row mass option_table_0019 seed_terms, both log-on causes q = 1/3, C2 silent; held out = impact after 00:19:37.443 imposed; D4/D5 Holland transient bias added. Uncontrolled-arm dynamics not calibrated (Boeing leave-one-out gate failed); in-flight structural outcome and pitch-down PROVISIONAL. (a) Kish row ESS of m x P; parents = distinct hand-off parents; run J for both-BFO cells (3rd and 4th) from the both-burst supplement (8 seeds), else production. (b) impact point over all rows; seabed = settling run-j-a1branch (GLORYS12V1, ERA5, TEOS-10, AusSeabed/GEBCO_2026; breakup rows settled from the intact impact point; afloat pieces excluded), field 'mean', Z = point Z + paired difference. (c) settled. (d) per-cell model (q from ATSB 2017 Fig. 73 lower-confidence fraction, rho terrain-weighted, OI density from AIS); grey = run J spatial (four-strata mixture). Drift, hydroacoustic and other evidence not used.