
Where the wreckage is on the seabed, for each use of the 00:19 satellite data

Flight families at the fixed weights of the flight model; impacts faster than 212 m/s removed (the model has no structural speed limit)



Summary (STE): Grey areas show where the wreckage is on the seabed. Dashed orange lines show where the aircraft hit the water. The movement of the wreckage through the water adds less than 1 % to each grey area. The flight model is not yet stable, so these areas can change. There are not yet sufficient samples to show a result for Holland H1 and H2.

Technical: Settled-mass seabed density (real ocean: GLORYS12V1 column 75-115 E, 45-10 S, TEOS-10 density; GLORYS12V1 surface current; ERA5 wind; AusSeabed then GEBCO_2026), settling 9330d4c, breakup table PROVISIONAL, afloat pieces excluded. Impacts: end of flight on core run C (4 strata x 8 seeds, prior track 289.7 deg, EoF 87f6343), option_posteriors with +unpowered (airborne at 00:19:37, not powered at 01:15:56). Strata weighted by core's probability of each flight family only (fixed weights); P(family) for R600 BTO + Raw BFO = 0.590 / 0.252 / 0.135 / 0.024 (free / repro-radar / descent-climb / routes). Log-on cause: other (H1: fuel exhaustion). Systematic resample, one settling draw each: 200,000 (a), 40,000 (b-e). 0.02 deg grid, Gaussian 0.1 deg, HPD. Mixture ESS by Kish; below 1,000 not estimable (H1/H2: descent-model and sampler changes await Pete; architecture burst study). Labels: core run C split-half NOT converged (min 0.82-0.86 against floor 0.924); rapid descents above 6,500 ft/min and unloading not reachable by the model; residual roll P(left) = 0.8 PROVISIONAL; trim at loss and two-tank takeover NOT applied; Mac, 2 threads. Infeasible-speed rows removed (model has no structural limit): rows with contact speed above 212 m/s (1.25 VMO) are removed and the rest re-weighted (architecture ruling 20:35 -0600 10 Oct, G-H2); removed weight 9.8-11.9 % (Held Out, R600), 44-47 % (H1, H2).