

Note (STE100): This chart shows where the aircraft can be on the sea floor. It uses the satellite data, flight models, debris drift, the searched areas and the Australian coasts where no debris was found. Only the coast near the cities and inner regional towns counts, at the same level as the coasts where debris was found. All other Australian coast is not used. This Australian part is a separate case, not the main result. There are few samples north of 33 S. Dark grey: 50 % of the probability; mid grey: 90 %; light grey: 99 %. Red line: 90 % without drift, searches and Australia.
Technical: Australian non-finds MEASURED by ocean drift, populated coast only (column In_I_lev_populated: ABS major city + inner regional coast at the find-coast level, outer regional and remote coast off; Pete's ruling 15 Oct 2026), dedicated arc-latitude run 20-40 S, both ocean models. core run J (tag 1244b7c; split-half accepted by ruling 16 Oct 2026); prior track = four-stratum mixture (waypoint routes 0.998, free flight, descent-climb, Davey dynamics with radar); waypoint routes with the run J northern supplement (rule B), stratified at 33 S; log-on cause: competing causes, prior odds 1/3, other cause silent; end-of-flight uncontrolled arm not calibrated; settling run J (breakup table provisional); seabed = settled pieces. Searched areas: per-cell detection model V1 central, Ocean Infinity coverage inferred (grade C), included as best estimate. Drift: finds, arc-latitude profile 20-40 S, GLORYS12 + GlobCurrent marginalised jointly. Hydroacoustics zero weight by ruling; Pleiades not used. Australian non-finds are a LABELLED SENSITIVITY for the decision table (Pete, 21 Oct 2026), not the headline. North of 33 S is under-sampled (effective sample from the seed scatter 168-341 for these two products, below the 1,000 target; table). HPD regions on the 769 km2 equal-area grid; planned next Ocean Infinity area is a map layer only (not drawn).