Effective sample size of each combination of 00:19 data, log-on cause and evidence



(i) Each number is the effective number of impacts that carry the result. Below 1,000 (red box), the result is not reliable. When we add the R1200 frequency, the drift and the Pléiades objects, few impacts remain. The fuel-exhaustion family without control (right) is the weakest.
(ii) Technical: Core run I: 4 core families x 8 seeds x 100,000 hand-off rows, equal family prior; NOT CONVERGED (split-half waypoints 0.916, floor 0.924; free flight 0.830, descent-climb 0.794, Davey reproduction 0.891). End of flight run I (binary mh370-eof-ri1, 102.4 M impacts): piloted turns and interim pitch-down provisional; no structural limit (G13). 00:19: competing log-on causes q = 1/3, W = 00:10:58-01:15:56, C2 silent, survival factor applied; no treatment uses the R1200 BTO. Ocean: GLORYS12 and GlobCurrent (product-relative windage), equal prior, mixed per row (Bayesian model averaging). Drift h50; unscored rows carried at the mean ratio (<= 0.11 % of weight); Western Australia non-recovery not modelled (G26). Hydroacoustics predictive only, not in these products. Searches: ATSB Phase 2 + Bluefin-21, rho 0.05, point target; Ocean Infinity excluded. Pléiades: rho 0.25, rating 5, six clusters plus four COSMO-SkyMed contacts, conditional on the objects being debris. Strata mixed by core share x 00:19 evidence (ruling C), conditional on equal family prior. Composer pass 2, architecture stand-in. Rows: 00:19 data treatment D0-D5 x log-on cause (q = 1/3 primary; C1, C2 silent, q = 1 sensitivities). Holland's Hypothesis 1 = D4 x C1, Hypothesis 2 = D3 x C2 silent. ESS on the equal-prior core mixture (ruling C), pooled over 8 seeds, ocean models mixed.