


Note (STE100): This chart shows where the aircraft can be on the sea floor, from the evidence that we use. Satellite data, flight models, debris drift and the searched areas give the result. The searched areas remove probability where the search was good. We do not use the hydrophone data. The Pleiades objects and the Australian coast are given only as separate cases. Top: the chance that the search did not find a wreck that was there, at 0, 1, 5 and 10 times the central value. Bottom: four levels of search quality; the open squares show two other ways to model the Ocean Infinity searches.

Technical: core run J (tag 1244b7c; split-half accepted by ruling 16 Oct 2026), prior track = four-stratum mixture (waypoint routes 0.998, free flight, descent-climb, Davey dynamics with radar); log-on cause: competing causes, prior odds 1/3, other cause silent; end-of-flight uncontrolled arm not calibrated; settling run J (breakup table provisional); seabed = settled pieces. Searched areas: per-cell detection model V1 central (q 0.95 / 0.70 by ATSB Fig. 73 lower-confidence fraction, rho = 0.05 x terrain weight, cap 0.95), Ocean Infinity coverage inferred (grade C), included as best estimate. Drift: finds only, dedicated arc-latitude run 20-40 S (run A outside), GLORYS12 + GlobCurrent marginalised jointly. Hydroacoustics zero weight by ruling; Pleiades conditional; Australian non-recovery excluded (sensitivity); planned next Ocean Infinity area is a map layer only. North of 33 S includes the run J northern supplement; north of 33 S under-sampled (effective sample in the tables). Views v3.2: waypoint routes use the run J northern supplement (rule B; settling d0660ec2, searched areas a8e7e28), stratified at 33 S by impact latitude, with A1 under D3-D5 from end of flight's two conditional strata (end-of-flight share); drift arc-latitude profile extended to 20-40 S (181a5fa). D5 waypoints at BFO error 4.3177 Hz; other strata as before. rho0 = 0 / 0.05 / 0.25 / 0.5 with the same terrain weight (V1_rho0, V1_central, V1_rho0.25, V1_rho0.5); levels V3, V1, V0, V2 of the per-cell model (237090ab); OI rows: uniform coverage fractions 0.889 / 0.7808, and without OI (sensitivity rows only). Error bars: seed-bootstrap SE (400 draws).