
Most probable impact regions, primary case (00:19 R600 BTO Only)

Dark: 50 % region. Light: 90 % region. +: highest density. Blue outline on the lower maps: the 90 % region above.



NOT CONVERGED (core run C). The dark area holds half the probability, and the dark and light areas together hold 90 %. The lower maps are correct only if the Pléiades objects are aircraft debris. The sample did not settle. Do not use these maps for a search.

Technical: Source: core run C (prior track 289.7 deg, 00:11 hand-off), strata free, Davey dynamics + radar, descent-climb, routes x 8 seeds, ~3.2 M impact rows per seed; core split-half mean 0.850-0.918 against floor 0.924: NOT CONVERGED. End of flight run C (binary 15ba1098; dive class (b); descents <= 6,500 ft/min; no structural limit, G13; residual roll P(left) 0.8 PROVISIONAL); log-on cause 'other', constraint +unpowered. Drift: GLORYS12V1 + ERA5 only, current 367 nodes, 50 km bandwidth, 9 finds; GlobCurrent excluded (F1); not-computed rows carried. Hydroacoustics: stand-in L_hyd scenario A with audit F8 (Blackman App. B noise), PREDICTIVE, NOT CALIBRATED (C6). Pléiades v2: rating 5, equal clusters, GLORYS12, given H. Searches: Phase 2 + Bluefin-21, rho 0.05. Strata mixed by core P(family) re-weighted by the 00:19 evidence. Holland H1/H2 not estimable, not shown. Regions: highest-density cells of the patch-B equal-area grid (0.25 deg lon x equal authalic-latitude steps, 769.3 km² cells), unsmoothed; peak from the map smoothed by 1 cell. All impacts (no speed filter).