Pléiades agreement test and sample adequacy



NOT CONVERGED (core run C). Left: a value above zero shows that the drift and sound evidence moves the probability towards the Pléiades area. Small dots are the eight seeds. Right: the number of independent samples that each step keeps. The sample did not settle.

Technical: Source: core run C (prior track 289.7 deg, 00:11 hand-off), strata free, Davey dynamics + radar, descent-climb, routes x 8 seeds, ~3.2 M impact rows per seed; core split-half mean 0.850-0.918 against floor 0.924: NOT CONVERGED. End of flight run C (binary 15ba1098; dive class (b); descents <= 6,500 ft/min; no structural limit, G13; residual roll P(left) 0.8 PROVISIONAL); log-on cause 'other', constraint +unpowered. Drift: GLORYS12V1 + ERA5 only, current 367 nodes, 50 km bandwidth, 9 finds; GlobCurrent excluded (F1); not-computed rows carried. Hydroacoustics: stand-in L_hyd scenario A with audit F8 (Blackman App. B noise), PREDICTIVE, NOT CALIBRATED (C6). Pléiades v2: rating 5, equal clusters, GLORYS12, given H. Searches: Phase 2 + Bluefin-21, rho 0.05. Strata mixed by core P(family) re-weighted by the 00:19 evidence. Holland H1/H2 not estimable, not shown. ln R = ln E[L_P | flight+EoF+drift+hydro] - ln E[L_P | flight+EoF] (Pléiades A_scene cancels); mixture over 00:19-re-weighted strata. Triangle: P + four COSMO contacts with correlated transport errors, rho 0.5 (joint-surface-rho0.5). ESS: composer Ess.parents.