

Simple: Drift run at 84 points along the 7th arc, 20–40° S (18 new points at 20–25° S on 20 Oct, 66 points at 26–40° S from 18 Oct). Each latitude has a point on the most likely impact track and points 50 NM either side. Error bars are 2 standard deviations of Monte Carlo noise from split halves; a repeat with new random seeds at 26° S shows that the true noise is about 1.5 times larger. Few flight paths end north of 30° S, so the shape of the PDF there is not reliable. Finds only is the base setting; the Australian no-find settings are labelled sensitivities.

Technical: Drift module, aus160 configuration (15–160 E), binary cbe08758, 3e5 particles per point at 20–32° S (1° steps), 1e6 at 33–40° S (0.5° steps); source grid extended to 19.0° S (margin 100 to 280 NM, same 5 NM lattice). Recovery bandwidth 50 km, window 8 Mar 2014–30 Jun 2016, 3 object classes, 4 ocean-error realisations, equal-weight mixture in likelihood of GLORYS12+ERA5 and GlobCurrent+ERA5 (product-relative windage). Detection relative to find coasts: option B 1/0.3/0.05 (city+inner regional/outer regional/remote, pre-registered), option C 0.3/0.1/0.02, populated only 1/0/0 (Pete's 15 Oct ruling); log-normal SD 1. Panel a: track points; 26° S repeat = same settings and seeds. Panel b: run J D0 impact histogram (00:19 data not used; all end-of-flight families: A1 cruise to fuel exhaustion then no control, A2 cruise to fuel exhaustion then controlled descent, B planned descent begun before fuel exhaustion), weighted by inverse-distance (6 nearest points) log-likelihood; 0.007% of flight-path mass lies outside 19.6–40.4° S. Davey et al. (2016), Fig. 10.3: recovered from the PDF vector paths (BTO+BFO). Northern supplement not included. Sensitivity, not a headline.