

Simple version (ASD-STE100): Each bar shows how much better the waypoint paths agree with one item of data than the other paths do. A bar to the right helps the waypoint paths. The largest gains are at the radar band between VAMPI and MEKAR (18:15 to 18:21), at the frequency data of the 18:39 phone call, and at the satellite data at 19:41. The waypoint paths fly exactly on the line from VAMPI to MEKAR, and the radar band is on that line. The total gain against free flight is 6.8. This gives the waypoint paths 99.8 % of the probability when the four groups have the same prior probability.

Technical version: Core run I, 8 seeds per family. Per data point, the stratum's sequential log-evidence increment ln p(y_k | y_1..k-1), modes combined by their prior weights; bars are waypoints minus the other family, mean over seeds, error bars the seed standard deviation (zero where all seeds share the same early particles). The increments sum exactly to the difference in total log-evidence; each is conditional on all earlier items, so the split depends on data order. Radar band = across-track scores against the VAMPI–MEKAR great circle at four times, sd 1.5 NM (radar-full.toml). The waypoint family flies the VAMPI–MEKAR prefix with no lateral randomness from 18:01:49 to 18:21; the other families carry lateral dispersion there. 18:39 and 23:15 are C-channel calls: BFO only, no BTO. 00:19: core, Davey both-BTO no-BFO.