
Paths that follow waypoints, which the data favour, turn south between IGOGU and SAMAK at 18:37–18:39 UTC



Simple version (ASD-STE100): The lines show possible flight paths from 18:22 to 18:42 UTC. Each line is one sample from the result. Blue lines are paths that fly to fixed waypoints. Orange lines are paths that fly freely. We show the same number of lines for the two groups, but the data give 99.8 % of the probability to the waypoint paths. The percentages show how many paths of each group turn in each area. The ± value is the change between the eight separate runs. The chart is for orientation only.

Technical version: Core run I, 12 Oct 2026; equal prior over four families (free flight, waypoints, descent-climb, Davey reproduction); family posterior from pooled log-evidence (waypoints −129.00, free −135.78). 8 seeds × 7M particles; 2,000 posterior route draws per seed, 40 s trace 18:22:00–18:42:40; 150 draws per seed per family drawn. Turn area classified by the northernmost latitude reached (≥7.55°N, 7.2–7.55°N, <7.2°N); ± is the seed standard deviation. Split-half = mean overlap over all 35 balanced seed partitions of the 00:19 latitude density (8-seed floor 0.924). Speed rule: early phase Mach 0.45–0.87 within 210–330 KCAS to 18:40, cruise 0.70–0.87 within 210–330 KCAS. 26 fixes (2014-epoch tiers A+B); fly-by and fly-over sampled; break-off 50/50 at 20/40/60/80 % on any leg. 00:19 treatment: core, Davey both-BTO no-BFO. Background: chart supplied by Pete (current cycle), affine fit to 12 fixes, residual ≤1.4 px; fix symbols at the positions used in the runs.