


Pléiades agreement test and sample adequacy

00:19 R600 BTO Only 00:19 Held Out



PIPELINE TEST - NOT CONVERGED (core (b) split-half). Left: a value above zero shows that the drift and sound evidence moves the probability towards the Pléiades area. Small dots are the four seeds. Right: the number of independent samples that each step keeps. The input sample did not settle.

Technical: Source: core (b) filter, prior track 289.7 deg, 00:11 hand-off, 4 strata x 4 seeds (51.2 M impact rows); end of flight 3c6319f (dive class (b), descent rates <= 6,500 ft/min, PROVISIONAL); log-on cause 'other', +alive. Drift: GLORYS12V1 + ERA5 only, 50 km bandwidth, 9 finds (GlobCurrent excluded: audit F1). Hydroacoustics: stand-in L_hyd (IMOS 4 loggers + Kadri H01W list). Pléiades: rating 5, 6 clusters, equal weight, GLORYS12, COSMO excluded. Searches: Phase 2 + Bluefin-21, rho 0.05, OI excluded. Not-computed rows carried at the (seed, mode) mean likelihood ratio (drift 16-25 % of weight, Pléiades 1-3 %). Strata mixed by P(family) re-weighted by the 00:19 evidence. Holland H1/H2 not estimable and not shown. In R = ln E[L_P | flight+EoF+drift+hydro] - ln E[L_P | flight+EoF] (the Pléiades A_scene constant cancels); mixture uses the 00:19-re-weighted strata. ESS: composer Ess.parents (children of one hand-off row summed).