


● Flight and descent only ● Add debris drift and underwater sound ● Also add the Pléiades objects (if aircraft debris)

Impact latitude for each end-of-flight family, primary case



PIPELINE TEST - NOT CONVERGED (core (b) split-half). Each line shows the impact latitude for one type of end of flight. P is the probability of that type. The data after 00:11 do not much change these probabilities, so they come mostly from the prior. Type B is only partly modelled.

Technical: Source: core (b) filter, prior track 289.7 deg, 00:11 hand-off, 4 strata x 4 seeds (51.2 M impact rows); end of flight 3c6319f (dive class (b), descent rates $\leq 6,500$ ft/min, PROVISIONAL); log-on cause 'other', +alive. Drift: GLORYS12V1 + ERA5 only, 50 km bandwidth, 9 finds (GlobCurrent excluded: audit F1). Hydroacoustics: stand-in L_hyd (IMOS 4 loggers + Kadri H01W list). Pléiades: rating 5, 6 clusters, equal weight, GLORYS12, COSMO excluded. Searches: Phase 2 + Bluefin-21, ρ 0.05, OI excluded. Not-computed rows carried at the (seed, mode) mean likelihood ratio (drift 16-25 % of weight, Pléiades 1-3 %). Strata mixed by P(family) re-weighted by the 00:19 evidence. Holland H1/H2 not estimable and not shown. Families from EoF family_labels (onset_mechanism x control_realised; 4 'A, controlled then lost' composed with A1, sensitivity with A2 in families.csv). B covers only descents begun after 00:11.