

IN SIMPLE WORDS. For each 00:19 data option, the left panel shows the probability that is left after the seabed searches, which found nothing. The right panel shows the area that holds 90 per cent of the probability after the searches. Four versions are shown: with all impacts or only impacts at 212 m/s or less, and with the groups of flight paths weighted by the core model only or also by the 00:19 data. Open markers are not estimable. All values are provisional.

TECHNICAL. run C NOT CONVERGED (free split-half 0.918 vs 0.924). Source: end of flight's run C sweep (4 strata x 8 seeds, 102,400,064 impacts) on core run C 00:11 hand-offs. Core P(family) passed in: free 0.589, Davey dynamics + radar 0.252, descent-climb 0.135, routes 0.024. Search likelihood: module run.toml (ATSB Phase 2 + Bluefin-21, detection probability 0.945 / 0.900, undetectable fraction rho 0.05, point target, Ocean Infinity excluded). 00:19 options with end of flight's +alive constraint (Holland H1: start-up offset, fuel-exhaustion log-on). End-of-flight labels: rapid descents above 6,500 ft/min and unloading not reachable by the model; residual roll P(left) 0.8 PROVISIONAL; trim at loss and two-tank takeover not applied. Compact impacts rebuilt for mh370 evaluate by the Pleiades stub (position only). No wreckage-field coverage (no run C settling samples). Architecture stand-in run; module to review. Z = sum over impacts of posterior weight x P(no find); mixture Z linear in stratum weights. Fixed, <=212: P_core x kept fraction. Re-weighted: P_core x Zhat_00:19 (end of flight's per_seed; matches the published run C family-evidence ln Zhat for 3 strata, routes not published).