
98 % of floating debris from the bearing-arc crossing is 190-1,140 km from the Haixun 01 contact at 03:15 UTC on 5 April 2014



Plain: We put 100,000 floating objects at the point where the 306.18 deg bearing meets the 7th arc, and at points 50-200 km inside the arc, on 8 Mar 2014 at the impact time from the sound. The shading shows where the objects are at 03:15 UTC on 5 Apr 2014, when white objects were seen about 90 km from the Haixun 01 contact. Almost no objects come within 110 km of the contact. Two ocean models are shown. This is a forward drift only; it does not use the flight-path calculation.

Technical: debris-drift engine snapshot harness (drift d51d80a, untracked stand-in test `snapshots`), aus160 configs with 28-day forcing (GLORYS12V1 0.494 m or GlobCurrent 0 m, ERA5 U10; GlobCurrent product-relative windage U(-0.75,-0.60) % U10), 3 object classes x 4 ocean-error/diffusivity realisations (K log-uniform 100-1000 m2/s), 8,334 per class, refloat off, 6 h step. Release at branch T impact times (00:31:52-00:34:24 UTC; branch C +2 min 27 s gives the same map within Monte Carlo noise). Positions linearly interpolated 00Z-12Z 5 Apr. HPD of object density, 0.1 deg bins. No flight-path posterior, no 00:19 option, no search case, no exclusions. Sighting: JACC 5-6 Apr 2014 (grade A), position not published.