


STE100: These maps show how well the 2018 seabed search north of 33° S could find the wreck at each place. (a) uses the positions of the search ships. (b) uses the shape of the seabed where it was mapped in 2014–2017. (c) shows where the remaining probability goes when these maps replace one fixed value. Blue: less probability. Red: more probability. North of 33° S the change is small, because little probability is there.
Technical: Ocean Infinity 2018 ground outside ATSB Phase 2, 0.005° grid (grid ends 24.5° S; Bluefin-21 at 21° S not shown; no 2025–26 ground north of 33° S). (a) d = 0.9·c, c = coverage density from slow AIS fixes (< 4 kn, MH370-CAPTION hourly), Gaussian σ 15 km, × traced layer, scaled to 112,000 km², capped at the layer (grade C, inferred). (b) ρ = 0.05·w, w = terrain weight (GA Phase 1 150 m bathymetry), mean 1 on Phase 2; w = 1 where there is no bathymetry (7–82% of ground by latitude band, 33–24.5° S). (c) Core run J × seabed searches, 00:19 R600 BTO + both BFOs as recorded (D3), settled wreckage, both-burst waypoints set (40,000 rows), per-cell model V1 minus scalar base V0 (OI uniform 0.889, q 0.9, ρ 0.05 flat), 0.2° bins, impact positions; D3 share north of 33° S: 0.015 before the searches, 0.044 after (V0), 0.041 after (V1). Fields and scores as corrected 20 Oct 2026 (w fallback and off-grid w). Core run J accepted as a pass by project ruling, 16 Oct 2026.