


(i) We used the sound measured at Wake Island and our own sound-loss model on the same path to find how much of the crash energy became sound. The answer depends on how deep the source was and on how long it lasted. For a source longer than about 1 s the answer stays the same. These are estimates for one crash of a small, fast jet.

(ii) eta_A = SE_obs / [E rho c/(2 pi) sum_b w_b S_b(tau) 10^(-TL_b/10)]; SE_obs = noise-subtracted exposure from Metz Fig. 3C/D (H11N 0.270, H11S 0.090 Pa^2 s); E = 900 MJ (Brown et al. 2026); TL = KRAKEN adiabatic on GEBCO_2026 + WOA23 B5C2 April, 5-63 Hz, plus the module's RAM correction; one-pole source, f^-2. Anchor (pre-registered, prereg 7dc315f): z_s 10 m, geometric mean of the triads = 2.26e-03 (tau_5 = 10 s at the grid edge, not identified; same value at the fallback 3.30 s). Spread: triads 3.3e-03/1.5e-03; E 700/1100 MJ 2.9e-03/1.8e-03; f^-4 1.3e-01; KRAKEN only 1.0e-03; +/-8 km 2.2e-03/2.4e-03. Measured-spectrum route: in-band 4-64 Hz coupling (lower bound on the total). Caveats C1 (propagation slope) and C2 (coupling not calibrated for an airliner) apply. All F-35A data DIGITISED from Metz, Obana & Fukao (2023, Pure Appl. Geophys. 180:1343).