



Predicted SOFAR arrivals at the IMS stations against the spans of Kadri's published traces

— +alive (reference) — 5-95 % ○ median Kadri (2024) Fig. 9 panel span
— +silent (sensitivity) — 0.5-99.5 % ► tail unconverged (> 5 min seed spread)



Run: reference-289 end-of-flight impacts (runs/snap289-m0011 hand-off; exchange eof-289-full seeds 1-4, 3.2M rows each), prior track $289.7^\circ \pm 1.0^\circ$, configs davey2016.toml + sensitivity/no-exhaustion-prior.toml + sensitivity/reference-snapshots.toml, rev 55c4536-dirty; Mac arm64. Uncorrected fuel; provisional sampler; PROVISIONAL-OVERNIGHT.
 Weights: end of flight's option_posteriors (displacement_hist.py @15ba915, read-only), option x log-on cause; +alive = airborne at 00:19:37.443 (EoF provisional reference), +silent = +alive, unpowered at 01:15:56, no later APU log-on (sensitivity). Seeds pooled equally. Validation vs EoF shares passed ($|\Delta\text{share}| \leq 1e-17$, $|\Delta q| \leq 1$ s).
 Arrival = impact time + WGS84 geodesic / c, c ~ N(1.482, 0.006) km/s (SOFAR); AGW c ~ U(1.40, 1.50) km/s (assumed, not modelled). No refraction or blockage path lengthening. Rows in (a,b): arms with pooled ESS $\geq 1,000$, ordered by +alive ESS. Grey: Kadri (2024) Fig. 9 panel spans. Script prepare/search_windows.py, prereq 55eb191.