



Predicted SOFAR arrivals from core (b) impacts, P(family) mixture, against reference-289



Run: hydroacoustics search windows, run by an architecture stand-in on the module's behalf; module to review. Labels: core (b): split-half NOT converged; two-tank bookkeeping only; PROVISIONAL-OVERNIGHT; EoF sweep run by a stand-in; uncorrected fuel; provisional sampler.

Impacts: mih370-exchange/end-of-flight/next-run (EoF 3c6319f, binary bcb6b252...), 4 strata x 4 seeds, ~3.2M rows/seed, from core (b) m0011 hand-offs; descent idle floor ON (off in reference-289); core s6-tanks dropped; single fuel pool, no one-engine phase. Prior track 289.7 ± 1.0 deg, deskstar, Innarsat ephemeris, internal-v1 fuel.

Strata combined by core's P(family), held fixed and unconverged: free 0.6948, Davey dynamics + radar 0.1527, descent-climb 0.1376, routes 0.0149; seeds equal-weight within stratum. Replicate k = seed k of every stratum mixed by P(family); convergence rule applied to the 4 replicates.

Method: prepare/search_windows.py @ 438c9e9 (sha256 1b05492b..., unchanged), weights from EoF option_posteriors (displacement_hist.py @ 15ba915, sha256 f964a9b9...), SATCOM sha256 8ec8c211...; stand-in mixture wrapper standin_mixture.py reproduces the script per stratum exactly (0 of 4,896 quantiles differ).

Arrival: WGS84 geodesic / c, c ~ N(1.482, 0.006) km/s (SOFAR); AGW c ~ U(1.40, 1.50) km/s, an allowance not a model; no refraction or blockage lengthening. Window = [q0.5 - 10 min, q99.5 + 20 min], 5-min rounding, widened to replicate envelope if spread > 5 min; union over 20 arms with mixture Kish ESS ≥ 1,000.

Variants: +alive = airborne at 00:19:37.443; +silent = +alive and unpowered at 01:15:56 (EoF 04:05 UTC 10 Oct, provisional). Arms ordered by mixture ESS. Holland H1/H2 (both_startup-offset-fuel-exhaustion, ESS 86; both_no-offset-other, ESS 124) not estimable, not shown. Reference-289: module run 438c9e9 on eof-289-full (snap289-m0011).