Fuel exhaustion then no control, under the satellite-object hypothesis: the branch pass raises the effective impacts in 8 of 9 cells (2 to 11 times), but no cell reaches the gate of 1,000



Footnote (simple): The data is conditional on the hypothesis that one or more objects seen by satellite on 20-23 March 2014 came from the aircraft. We do not use the shape of the objects. The end of flight physics is provisional. No seabed search is applied. The aircraft cruises to fuel exhaustion, then has no control. 'Run J sample' is the production sample; 'with branch pass' adds 8 seeds that sample only the flame-out, no-control descents. The search gate is 1,000 effective impacts.
Footnote (technical): Core run J (split-half accepted as a pass by project ruling, 16 Oct 2026), waypoints stratum (P 0.998), 24 production seeds; D3/D4 production = both-burst set (8 seeds, thinned). A1-branch conditional pass: 8 seeds x 100k hand-offs, branch = onset flame-out-associated and control realised none; stratified estimator Z(d,B) = Z_prod(D0',c,B) E_pass[L_d L_H]/E_pass[L_D0'] (branch replaced). H: Pleiades rating 5 (6 clusters) + COSMO-SkyMed F1-F4, pass-marginal; rho 0.25; GLORYS12+ERA5 and GlobCurrent+ERA5 (product-relative windage), equal prior. Kish ESS of A1 rows (pass rows scaled by alpha plus production A1 outside the branch). No identity likelihood; no Bayes factor or P(H). Before any seabed search.