

Notes. Core: satellite communication and flight-performance posterior of end-of-flight run J, with the 00:19 data held out. Drift: two ocean models with equal weight (GLORYS12V1 currents at 0.494 m with ERA5 wind; Copernicus GlobCurrent surface currents with ERA5 wind; the GlobCurrent wind factor is reduced by 0.60 to 0.75 per cent for its built-in wind drift). Flaperon response as measured by CSIRO (1.2 per cent wind factor, 0.10 m/s leeway); other exterior panels 1.2 +/- 0.3 per cent; interior panels median 2.5 per cent. 84 release points along the 7th arc; 6250 particles per object type and per ocean sample; 16 ocean samples, 32 south of 33.25 S. Finds scored to 30 Jan 2017 within 50 km of the find place; delay from arrival to find exponential, mean 60 days. Detection levels log-normal (sigma 1) per coast and period, 16384 draws; prior median 0.1 of the full level before 29 Jul 2015 (flaperon find), 0.5 to 1 Mar 2016, then 1. Coasts: Reunion, Mauritius and Rodrigues, Mozambique, South Africa, Madagascar, Pemba and the rest of East Africa, Comoros, Seychelles. Coasts with few people or little searching start at 0.3 of the level of the find coasts. Australian coasts carry no information unless stated. Twelve items: the nine finds of Durgadoo et al. and Linga Linga, Nautilus Bay and Mpame, weighted by their identification grade (confirmed 1, almost certain 0.98, highly likely 0.9). Not used: items identified partly from drift (sensitivity only), Sandravinany (date range past the window), coasts in Asia. Provisional physics; Monte Carlo uncertainty of the mean about 0.2 degrees.