
Model changes, one at a time (nine finds unless stated)
As first scored: to 30 Jan 2017, three periods, six find coasts
Window to 30 Jun 2016 (as on 7 Nov), this run
Fourth detection period from 1 Jul 2016
All western Indian Ocean coasts
All coasts and four periods
Headline: twelve items, weighted by identification grade
Changes to the headline
All identification weights set to 1
Add eight items identified partly from drift
Without Nautilus Bay and Mpame
Fresh arrival: flaperon and Mossel Bay 10 days, Paindane 14 days
Fresh arrival: flaperon and Mossel Bay 3 days, Paindane 14 days
Mossel Bay find dated at its second beaching (Mar 2016)
Official alternative dates (Vilanculos, Pemba, Mauritius)
Detection level the same in all periods (no increase after the flaperon find)
Other coasts at the level of the find coasts
Other coasts at 0.1 of the level of the find coasts
Populated new South African and Mozambique coasts at 0.3
No finds on Australian coasts, ratio 1
No finds on Australian coasts, ratio 0.25
No finds on Australian coasts, ratio 0.05
All items except Mossel Bay
All items except Mossel Bay and Paindane
Mossel Bay only
Flaperon only
Find-place tolerance 25 km
Find-place tolerance 100 km
Find-place tolerance 200 km
GlobCurrent ocean model only
GLORYS12 ocean model only
0
1
2
Change in mean latitude (degrees north)