



Seabed search evidence against p , for three uses of the 00:19 data — NOT CONVERGED

The searches remove less probability when p is larger



p controls how much probability stays on searched ground



Plain (ASD-STE100): The chart shows how the result of the seabed searches changes with p , the probability that the wreck is in a searched area but the search did not find it. The impacts come from the core run (b) with the end-of-flight stage, four flight-path groups, mixed with weights updated by the 00:19 data (dashed lines: fixed weights). The aircraft was transmitting at 00:19:37 and had no electrical power at 01:15:56. The core run is not converged, so these values are provisional. No published value of p exists. The reference is 0.05. Davey et al. (2016, p. 102) use p close to 0.

Technical: $Z = \sum f P_f Z_f$; thick: P_f re-weighted by EoF $\hat{Z}_{00:19}$ (family-evidence-next-run-b.json, +alive keys); dashed: fixed $P = 0.697/0.152/0.141/0.010$ (module convention 0.69/0.15/0.14/0.01 renormalised). EoF next-run on core (b) m0011 hand-offs, 4 strata $\times$ 4 seeds $\times$ 3.2M = 51,200,096 impacts; weights from EoF option_posteriors, log-on cause other, +unpowered (airborne 00:19:37.443, not powered 01:15:56). Campaigns: Phase 2 (q 0.945, measured valid-data coverage) + Bluefin-21 (q 0.9); no Ocean Infinity. 'On searched ground' = $\sum w \cdot c$ Phase2. Holland H1/H2 not yet estimable - targeted sampler in progress (mixture ESS 85/123), not shown. Coverage gap: EoF descent rates above 6,500 ft/min and unloading not reachable by the model. UNCONVERGED: core (b) split-half NOT converged; two-tank bookkeeping only; right-dry rows twin-engine continuation, left pool drained at doubled grid_inop (internal-v1); PROVISIONAL-OVERNIGHT; EoF sweep and this analysis run by architecture stand-ins, searched areas to review.