

Where the wreckage is on the seabed, for each set of evidence

Grey: 90 % region of the wreckage. Colour: 50 % region. Dashed line: 90 % region of the impact point.

00:19 R600 BTO Only (primary)



00:19 Held Out



NOT CONVERGED (core run C). The areas show where the wreckage is possibly on the seabed. The right half is correct only if the Pléiades objects are aircraft debris. The wreckage moves very little as it sinks, so the seabed areas are almost the same as the impact areas. Do not use these maps for a search.

Technical: G panels: settling's published aimed samples (settling/next-run-c-composed-aimed); Pléiades panels: the stand-in's run of the same tools. Settling's aimed resample (results/settling-aimed-resample/wf_aimed.py) of 40,000 impacts drawn from the composed per-row weights (composer pass 1, run C, +unpowered; strata by 00:19-re-weighted P(family)); one settling draw each with settling's wreckage_field test (copy of settling 9330d4c; real ocean GLORYS12V1 column, breakup table PROVISIONAL). After searches = composed weight x searched-areas likelihood per row (point target, rho 0.05). Factors: EoF run C (no structural limit, G13; descents <= 6,500 ft/min), drift GLORYS12 only, current nodes (GlobCurrent excluded, F1; not-computed rows carried), stand-in hydro + F8 (predictive, not calibrated), Pléiades v2 rating 5, equal, GLORYS12, given H. 0.02 deg grid, Gaussian 0.1 deg, HPD on settled mass; afloat pieces excluded. Row ESS (Kish) >= 1,000 required. Holland H1/H2 not estimable.