




Outliers from the H08S tests on Kadri (2024) Fig. 9 vectors (filtered, single channel; not raw IMS): cycle test h08s_shot_outliers.py (ddaa848/4b0ee7e), energy-only test h08s_pulse_energy.py (8ff37fb) and the exploratory off-cadence search. Each is paired with every H01W trigger (2a detector, strict or loose) or bearing-gated Kadri Table 1 event whose H08S–H01W time difference lies in 856–1,347 s. The implied source and impact time are prior-weighted means using both arrival times; impact 00:19:37 + 300 $\pm$ 180 s, $c = 1.482 \pm 0.006$ km/s, 10 s picks; Table 1 bearings with Student- t sd 3.3°. Filled = $\log_{10}$ BF ≥ 1 against a chance pair; open = below; bars = ± 1 sd. Every pre-registered count is consistent with chance ($p \geq 0.20$). The grey map prior and the grey impact-time curve are the stand-in from run no-exhaustion-prior (b3dd44b), prior track $295.66^\circ \pm 1.0^\circ$, config sensitivity/no-exhaustion-prior.toml, α 20 NM; the scoring uses this stand-in. The blue curves are the reference-289 impact times from eof-289-full seed 1 (source run snap289-m0011, prior track $289.7^\circ \pm 1.0^\circ$, held-out 00:19 option, weight $\times \exp(\loglik:none)$; all, dotted = flight-path angle $< 10^\circ$, red dashed = $\geq 10^\circ$), shown for comparison only and not used in the scoring: 5–95 % -2.9 to $+46.3$ min, median 19.6 min, 0.7 % beyond the axis (tail to about 102 min). EXPLORATORY characterisation.