


Seabed wreckage PDF under the four 00:19 priority hypotheses (reference-289, +alive (PROVISIONAL-OVERNIGHT))



(c) Holland H1: both bursts, start-up offset; fuel-exh.

**NOT ESTIMABLE
impact ESS 36 (< 1,000)**

90 % area, thousand km²: seabed 43, same impacts 41
40,000 draws on 281 distinct impacts; impact ESS 36

(d) Holland H2: both bursts, no offset; other

**NOT ESTIMABLE
impact ESS 82 (< 1,000)**

90 % area, thousand km²: seabed 59, same impacts 58
40,000 draws on 427 distinct impacts; impact ESS 82

seabed wreckage, 50 %

seabed wreckage, 99 %

6th arc, 00:11 UTC

seabed wreckage, 90 %

impact PDF 50/90/99 % (reference)

7th arc, 00:19 UTC

Grey: seabed wreckage PDF, weighted by settled mass (afloat pieces have no seabed position and are excluded). Dashed orange: impact PDF, every impact at full weight. Before any seabed-search evidence. Impacts: end of flight eof-289-full-s{1,2,3,4} (reference-289, prior track 289.7°, source runs/snap289-m0011), seeds pooled with equal weight; weights = option_posteriors (hand-off × 00:19 burst likelihood, × log-on lag density for fuel-exhaustion). Dive class and Boeing-calibrated glide PROVISIONAL-OVERNIGHT. Settling (hypothesis/settling 9823b4e): real ocean, GLORYS12V1 column (TEOS-10 density) and surface current, ERA5 wind, AusSeabed then GEBCO_2026; breakup table PROVISIONAL; implosion and occupants alternatives off. Systematic resampling, one settling draw per resampled impact (a: 179,558 of 200,000 kept by +alive; b: 39,999 of 40,000; c, d: 40,000). Impacts north of 18° S lie outside the ocean window: not computed, excluded (counted in the JSON). 0.02° grid, Gaussian 0.1° (6 NM), HPD on the smoothed density. NOT ESTIMABLE: pooled impact ESS below 1,000; such a panel shows where a few dozen impacts lie, not a posterior. Platform: Mac, 2 threads. Run labels: uncorrected fuel (fuel-model audit F1-F14 open); +alive: airborne at 00:19:37.443 (end of flight, PROVISIONAL-OVERNIGHT), applied by rejection to the plain resample.