


Seabed wreckage PDF under the four 00:19 priority hypotheses (reference-289)



Grey: seabed wreckage PDF, weighted by settled mass (afloat pieces have no seabed position and are excluded). Dashed orange: impact PDF, every impact at full weight. Before any seabed-search evidence. Impacts: end of flight eof-289-full-s{1,2,3,4} (reference-289, prior track 289.7°, source runs/snap289-m0011), seeds pooled with equal weight; weights = option_posteriors (hand-off × 00:19 burst likelihood, × log-on lag density for fuel-exhaustion). Dive class and Boeing-calibrated glide PROVISIONAL-OVERNIGHT. Settling (hypothesis/settling 9823b4e): real ocean, GLORYS12V1 column (TEOS-10 density) and surface current, ERA5 wind, AusSeabed then GEBCO_2026; breakup table PROVISIONAL; implosion and occupants alternatives off. Systematic resampling, one settling draw per resampled impact (200,000 in a; 40,000 in b-d). Impacts north of 18° S lie outside the ocean window: not computed, excluded (counted in the JSON). 0.02° grid, Gaussian 0.1° (6 NM), HPD on the smoothed density. NOT ESTIMABLE: pooled impact ESS below 1,000; such a panel shows where a few dozen impacts lie, not a posterior. Platform: Mac, 2 threads. Run labels: uncorrected fuel (fuel-model audit F1-F14 open).