Prior recovery: every corrected estimate agrees with the uncorrected run (all |z| below 1.6)

Quantity	Same population (run J seed 1)	Supplement, new seeds vs run J	New seeds, normal output vs run J	Combined (rule B) vs run J
Mean latitude	+0.76	-0.37	-0.40	+0.08
Mean longitude	+1.00	-0.47	-0.50	—
Mean altitude	-1.31	-0.51	-0.54	—
Mean Mach	-0.53	+0.56	+0.57	—
Mean BFO bias	-0.22	-0.45	-0.46	—
P(north of 33°S)	-0.00	+0.48	+0.49	+0.67
P(north of 30°S)	-0.15	+0.52	+0.48	+1.00
P(north of 25°S)	-0.34	+0.89	+0.82	+1.59
P(true heading)	+0.63	+0.54	+0.53	+0.58
P(magnetic heading)	+0.05	+0.58	+0.48	+0.67
P(true track)	-0.98	-0.66	-0.65	-0.39
P(magnetic track)	+1.35	+0.42	+0.40	+0.86
P(lateral navigation)	-0.13	+0.40	+0.41	-0.28
Median latitude at 00:19	—	—	+0.26	—
Mean latitude at 00:19	—	—	-0.77	—
P(north of 33°S) at 00:19	—	—	+0.05	—
P(north of 30°S) at 00:19	—	—	-0.36	—
P(34.5–36.5°S) at 00:19	—	—	-1.28	—

Plain: The table compares each corrected supplement estimate with the estimate that has no correction. A z value is the difference divided by its standard error. All values are less than 3 in size, so the correction does not change the result. Rows marked 00:19 use the final result, with both 00:19 BTO values and no 00:19 BFO. The source is the core filter: run J, early-flight family 'waypoints' (24 runs), and the northern supplement (16 new runs). The prior track starts from the 18:01 radar position and follows a random route through the 2014 waypoints. The 00:11 data are used; the 00:19 data are not. The one-engine physics and the fuel-flow factor are provisional. No search areas, ocean drift or end-of-flight data are applied. The 18:25:27 signal is not used. The other three early-flight families are not included.

Technical: Column 1: run J waypoints seed 1, supplement drawn from its own 00:11 population, against its normal hand-off; SE from Kish ESS over distinct states. Columns 2-3: new seeds 1001-1016 against run J seeds 1-24, seed-level z = difference of seed means / sqrt(s1^2/16 + s2^2/24). Column 4: rule B combination against run J alone, production seed SE only (conservative). 00:19 rows: final.npy of the new seeds (re-allocated modes) against run J. Pass criterion |z| < 3. Core 00:11 hand-offs (filtering distribution at m0011, look-ahead correction exp(log_correction) applied): run J waypoints seeds 1-24 (tag run-j, 1244b7c; 100k rows per seed) and supplement seeds 1001-1016 (rev 3bf0801; particles per mode [TH, MH, TT, MT, LN] = [1.2, 1.3, 1.1, 2.2, 1.2]M vs run J [1.0, 0.5, 2.5, 0.5, 2.5]M; supplement hand-off 100k rows per seed, tilt r(lat) = 0.1 + 0.9 h/H over 00:11 latitude -34.5..-21.0). BFO sd 4.3177 Hz; kappa_x ~ N(1.0, 0.046) and one-engine s7 + s8 PROVISIONAL; 18:25:27 R600 held out. Search case: none; ocean models: none; end-of-flight 00:19 options: not applied. Combination rule B: balance heuristic with r_hat (r_hat.csv), n1 = 2,399,999, n2 = 1,600,003 rows.